

Adversarial quantum channel discrimination

Kun Fang,^{1,*} Hamza Fawzi,^{2,†} and Omar Fawzi^{3,‡}

¹*School of Data Science, The Chinese University of Hong Kong, Shenzhen, Guangdong, 518172, China*

²*Department of Applied Mathematics and Theoretical Physics,*

University of Cambridge, Cambridge CB3 0WA, United Kingdom

³*Inria, ENS de Lyon, UCBL, LIP, 69342, Lyon Cedex 07, France*

(Dated: June 4, 2025)

We introduce a new framework for quantum channel discrimination in an adversarial setting, where the tester plays against an adversary who accesses the environmental system and possesses internal quantum memory to perform adaptive strategies. We show that in asymmetric hypothesis testing, the optimal type-II error exponent is precisely characterized by the minimum output channel divergence, a new notion of quantum channel divergence in the worst-case scenario. This serves as a direct analog of the quantum Stein’s lemma in the adversarial channel discrimination. Notably, the optimal error exponent can be achieved via simple non-adaptive strategies by the adversary, and its value can be efficiently computed despite its regularization. The strong converse property for quantum channel discrimination also holds in general. This adversarial quantum Stein’s lemma is proved by new chain rules for measured and sandwiched relative entropies. Moreover, we derive a generalized version of the entropy accumulation theorem between two arbitrary sequences of quantum channels, extending the existing results from entropy to divergence and providing a solution to the dual formulation of the open problem presented in [IEEE FOCS, pp. 844–850 (2022)].

A quantum channel describes the most general physical transformation that quantum states can undergo. It captures not only idealized processes like unitary evolutions but also non-ideal, noisy, or dissipative processes, making them crucial for understanding real-world quantum systems. The task of quantum channel discrimination is to identify which member of a given set of quantum channels governs the physical process in a black-box scenario. It is pivotal to various quantum information tasks [1–12] and gives insights into a wide range of quantum protocols and applications, from quantum foundations (e.g., exploring the quantum advantage of entanglement [13–16]) to quantum communication (e.g., estimating quantum channel capacity [17–22]), quantum sensing (e.g., quantum reading of classical data and quantum illumination of targets [23, 24]) and even quantum biology [25, 26].

In this work, we introduce an operationally-motivated adversarial framework for quantum channel discrimination, where a tester interacts with an *untrusted* quantum device that generates quantum states upon request. Such an adversarial setting arises naturally in other quantum information tasks such as quantum key distribution [27, 28], quantum adversarial learning [29], quantum interactive proofs [30] and quantum state verification [31], where ensuring reliable performance requires resilience against adversarial behavior. The task of adversarial discrimination was previously studied in the classical case by [32]. Here, we extend the investigation to the general quantum scenario [33]. Unlike the conventional channel discrimination model—where inputs are typically fixed or optimized by the tester to maximize distinguishability—the adversarial setting addresses the *worst-case* scenario where an unknown or malicious agent may manipulate the input to minimize the tester’s ability

to distinguish between channels. This can be seen as the problem of discriminating between two quantum devices by observing the output on an untrusted input.

Technically, the discrimination task is formulated as a hypothesis testing problem, with the goal of finding discrimination strategies that give the optimal trade-off between two kinds of error probabilities, namely the probabilities of false detection (type-I error) and false rejection (type-II error) [34, 35]. The challenge in adversarial discrimination arises from the potential capabilities of the adversary, who may access the channel’s environmental system—the ancillary quantum system that interacts with the main system and can cause information leakage—or possess (potentially unbounded) quantum memory that stores partial information from previous rounds and uses it in subsequent rounds to enable adaptive strategies. This raises a central question: how effectively can the tester distinguish between two quantum channels while playing against such an adversary?

This work provides a complete answer to the above question in the context of asymmetric hypothesis testing. Specifically, we show that in adversarial quantum channel discrimination, the optimal type-II error decays exponentially at a rate characterized by a new notion of quantum channel divergence, termed the minimum output channel divergence, provided the type-I error remains within a fixed threshold. This result serves as a direct analog of the quantum Stein’s lemma in the adversarial channel discrimination and it is proved by new chain rules for quantum relative entropies. Notably, the optimal error exponent can be achieved via simple non-adaptive strategies by the adversary, and desirable mathematical properties such as computational efficiency and the strong converse property hold in general. Extending

the adversarial discrimination framework and using the chain rules, we derive a relative entropy accumulation theorem, which states that the hypothesis testing relative entropy between two sequential processes of quantum channels can be lower bounded by the sum of the regularized minimum output channel divergences. A special case of this theorem recovers a version of the entropy accumulation theorem (a foundational tool in quantum cryptography [36–38]).

Minimum output channel divergence.— Let $\mathcal{D}(A)$, $\mathcal{L}(A)$ and $\mathcal{H}_+(A)$ denote the set of density operators, linear operators and positive semidefinite operators on a finite-dimensional Hilbert space \mathcal{H}_A , respectively. The set of completely positive maps from $\mathcal{L}(A)$ to $\mathcal{L}(B)$ is denoted by $\text{CP}(A : B)$. A quantum channel $\mathcal{N}_{A \rightarrow B}$ is a linear map from $\mathcal{L}(A)$ to $\mathcal{L}(B)$ that is both completely positive and trace-preserving. The set of all such maps is denoted by $\text{CPTP}(A : B)$. A quantum divergence is a functional $\mathbb{D} : \mathcal{D} \times \mathcal{H}_+ \rightarrow \mathbb{R}$ that satisfies the data-processing inequality, characterizing the “distinguishability” or “distance” between two quantum states. For any subsets $\mathcal{A} \subseteq \mathcal{D}$ and $\mathcal{B} \subseteq \mathcal{H}_+$, their divergence is defined as $\mathbb{D}(\mathcal{A} \parallel \mathcal{B}) := \inf \{ \mathbb{D}(\rho \parallel \sigma) : \rho \in \mathcal{A}, \sigma \in \mathcal{B} \}$, which represents the minimum “distance” between the sets.

Let $\mathcal{N} \in \text{CPTP}(A : B)$ and $\mathcal{M} \in \text{CP}(A : B)$. We introduce the *minimum output channel divergence* as

$$\mathbb{D}^{\text{inf}}(\mathcal{N} \parallel \mathcal{M}) := \inf \{ \mathbb{D}(\mathcal{N}(\rho) \parallel \mathcal{M}(\sigma)) : \rho, \sigma \in \mathcal{D}(A) \},$$

capturing the worst-case distinguishability of the channels under optimized test states. It is also useful to see this as a divergence between two sets of quantum states, $\mathbb{D}^{\text{inf}}(\mathcal{N} \parallel \mathcal{M}) = \mathbb{D}(\mathcal{L}(\mathcal{N}) \parallel \mathcal{L}(\mathcal{M}))$, where $\mathcal{L}(\mathcal{D}) := \{ \mathcal{L}(\rho) : \rho \in \mathcal{D} \}$ denotes the image set of \mathcal{D} under a linear map \mathcal{L} . The regularized channel divergence is defined as

$$\mathbb{D}^{\text{inf}, \infty}(\mathcal{N} \parallel \mathcal{M}) := \lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{D}^{\text{inf}}(\mathcal{N}^{\otimes n} \parallel \mathcal{M}^{\otimes n}),$$

which accounts for the asymptotic behavior of the channel divergence over multiple uses.

Adversarial quantum channel discrimination.— Consider a scenario where a tester interacts with an untrusted quantum device that generates quantum states upon request. The device guarantees that the states are produced by either a quantum channel \mathcal{N} or a quantum channel \mathcal{M} . The tester is allowed to request multiple samples from the device and perform measurements to distinguish between the two cases. More formally, let $\mathcal{N}_{A \rightarrow B}$ and $\mathcal{M}_{A \rightarrow B}$ be the two quantum channels to distinguish, and let $\mathcal{U}_{A \rightarrow BE}$ and $\mathcal{V}_{A \rightarrow BE}$ be their Stinespring dilations, respectively, with E denoting the environmental system. These channels satisfy the relations that $\mathcal{N}_{A \rightarrow B} = \text{Tr}_E \circ \mathcal{U}_{A \rightarrow BE}$ and $\mathcal{M}_{A \rightarrow B} = \text{Tr}_E \circ \mathcal{V}_{A \rightarrow BE}$, where $\mathcal{U}_{A \rightarrow BE}(\cdot) := U(\cdot)U^\dagger$ and $\mathcal{V}_{A \rightarrow BE}(\cdot) := V(\cdot)V^\dagger$, and Tr_E denotes the partial trace over the environment.




FIG. 1. Adaptive and nonadaptive strategies for two rounds of state generation, where \mathcal{U} and \mathcal{V} are the Stinespring dilations of quantum channels \mathcal{N} and \mathcal{M} , respectively, \mathcal{P}^i and \mathcal{Q}^i are internal operations by the adversary, I is the identity map, $\{M_2, I - M_2\}$ is a quantum measurement by the tester.

An *adaptive strategy* for adversarial quantum channel discrimination proceeds as follows (see Figure 1(a)). Suppose the device operates according to the channel \mathcal{N} . Initially, the adversary prepares a quantum state using a channel $\mathcal{P}^1 \in \text{CPTP}(R_0 E_0 : A_1 R_1)$, where $|R_0| = |E_0| = 1$, and sends system A_1 through the channel \mathcal{U} , generating the output state $\mathcal{U} \circ \mathcal{P}^1$ and returning the system B_1 to the tester. In the next round, the adversary performs an internal update $\mathcal{P}^2 \in \text{CPTP}(E_1 R_1 : A_2 R_2)$, leveraging information stored in the quantum memory R_1 and the environmental system E_1 from the previous round. The adversary sends system A_2 through the channel, producing the output state $\mathcal{U} \circ \mathcal{P}^2 \circ \mathcal{U} \circ \mathcal{P}^1$ and returning the system B_2 to the tester. This process is repeated for n rounds. Finally, the tester performs a quantum measurement $\{M_n, I - M_n\}$ on the collected systems $B_1 \cdots B_n$ in their possession to determine which channel was used to prepare the states.

After n rounds of state generation, the tester obtains an overall state on the systems $B_1 \cdots B_n$ as:

$$\rho[\{\mathcal{P}^i\}_{i=1}^n] := \text{Tr}_{R_n E_n} \prod_{i=1}^n \mathcal{U}_{A_i \rightarrow B_i E_i} \circ \mathcal{P}_{R_{i-1} E_{i-1} \rightarrow A_i R_i}^i.$$

Similarly, if the device is governed by \mathcal{M} and the internal operations by the adversary are given by \mathcal{Q}^i , then the overall state on system $B_1 \cdots B_n$ is given by

$$\sigma[\{\mathcal{Q}^i\}_{i=1}^n] := \text{Tr}_{R_n E_n} \prod_{i=1}^n \mathcal{V}_{A_i \rightarrow B_i E_i} \circ \mathcal{Q}_{R_{i-1} E_{i-1} \rightarrow A_i R_i}^i.$$

Due to limited knowledge of the internal workings of the device, the tester only has access to partial information, knowing that their state belongs to either: $\mathcal{A}_n :=$

$\{\rho[\{\mathcal{P}^i\}_{i=1}^n] : \mathcal{P}^i \in \text{CPTP}(R_{i-1}E_{i-1} : A_iR_i), \forall R_i, \forall i\}$ or $\mathcal{B}_n := \{\sigma[\{\mathcal{Q}^i\}_{i=1}^n] : \mathcal{Q}^i \in \text{CPTP}(R_{i-1}E_{i-1} : A_iR_i), \forall R_i, \forall i\}$ where the adversary's internal memory R_i can have arbitrarily large dimensions.

A *non-adaptive strategies* is a subclass of adversarial strategies that disregards the environmental systems E_i and performs no updates between rounds (see Figure 1(b)). Here, the operations $\mathcal{P}_i, \mathcal{Q}_i$ ($i \geq 2$) are simply identity maps, with $R_i = A_{i+1} \cdots A_n$. The sets of all possible outputs are denoted by \mathcal{A}'_n and \mathcal{B}'_n , respectively, which also correspond to the images of the tensor product channels, $\mathcal{A}'_n = \mathcal{N}^{\otimes n}(\mathcal{D})$ and $\mathcal{B}'_n = \mathcal{M}^{\otimes n}(\mathcal{D})$.

The adversarial quantum channel discrimination is essentially to distinguish between the two sets \mathcal{A}_n (\mathcal{A}'_n) and \mathcal{B}_n (\mathcal{B}'_n). Define the type-I error $\alpha(\mathcal{A}_n, M_n) := \sup_{\rho_n \in \mathcal{A}_n} \text{Tr}[\rho_n(I - M_n)]$ and type-II error $\beta(\mathcal{B}_n, M_n) := \sup_{\sigma_n \in \mathcal{B}_n} \text{Tr}[\sigma_n M_n]$. Asymmetric hypothesis testing investigates the decay of the optimal exponent of the type-II error probability when the type-I error is within a constant threshold ε , that is, to evaluate:

$$\beta_{n,\varepsilon}(\mathcal{N}||\mathcal{M}) := \inf_{0 \leq M_n \leq I} \{\beta(\mathcal{B}_n, M_n) : \alpha(\mathcal{A}_n, M_n) \leq \varepsilon\}.$$

Our main result, termed *adversarial quantum Stein's lemma* in Theorem 1, establishes that this optimal exponent is precisely characterized by the regularized minimum output channel divergence induced by the *Umegaki relative entropy* [39] $D(\rho||\sigma) := \text{Tr}[\rho(\log \rho - \log \sigma)]$ if $\text{supp}(\rho) \subseteq \text{supp}(\sigma)$ and $+\infty$ otherwise.

Theorem 1 (*Adversarial quantum Stein's lemma.*) *Let $\mathcal{N} \in \text{CPTP}(A : B)$ and $\mathcal{M} \in \text{CP}(A : B)$. Then for any $\varepsilon \in (0, 1)$, it holds that*

$$\lim_{n \rightarrow \infty} -\frac{1}{n} \log \beta_{n,\varepsilon}(\mathcal{N}||\mathcal{M}) = D^{\text{inf},\infty}(\mathcal{N}||\mathcal{M}), \quad (1)$$

where the Stein's exponent on the right-hand side can be efficiently computed via semidefinite programs. Moreover, the optimal exponent can be achieved by non-adaptive strategies, indicating that adaptive strategies by the adversary offer no advantage over non-adaptive ones.

The minimum output channel divergence $D^{\text{inf}}(\mathcal{N}||\mathcal{M})$ is not additive in general, necessitating the use of the regularized limit in the Stein's exponent $D^{\text{inf},\infty}(\mathcal{N}||\mathcal{M})$ [40]. While this could make its estimation challenging, the fact that the optimal exponent can be achieved by non-adaptive strategies simplifies the problem. As the sets \mathcal{A}'_n and \mathcal{B}'_n from non-adaptive strategies fall within the framework of the generalized asymptotic equipartition property (AEP) in [41], the Stein's exponent $D^{\text{inf},\infty}(\mathcal{N}||\mathcal{M})$ can be approximated within an additive error δ by a quantum relative entropy program of size $O((l+1)^k)$, where $k = \max\{|A|^2, |B|^2\}$ is given by the channel dimensions, and $l = \lceil \frac{8|B|^2}{\delta} \log \frac{|B|^2}{\delta} \rceil$ relates to the expected accuracy. Further details on this computational aspect can be found in the accompanying paper [40].

It is also worth noting that Eq. (1) established above universally applies to any $\varepsilon \in (0, 1)$, thereby demonstrating the ‘‘strong converse property’’, a desirable mathematical property in information theory [42] that delineates a sharp boundary for the tradeoff between the Type-I and Type-II errors in the asymptotic regime. That is, any discrimination strategy achieving a Type-II error decay rate exceeding the Stein's exponent $D^{\text{inf},\infty}(\mathcal{N}||\mathcal{M})$ will necessarily result in the Type-I error converging to one in the asymptotic limit. While this property has been proven for the discrimination between two quantum states [35], it remains an open question for discriminating two quantum channels in the best-case scenario [22]. Our result gives an answer in the worst-case scenario and contains quantum states [34, 35] as a special case when the channels are taken as replacer channels.

Chain rules and proof outline for Theorem 1.— The proof of Theorem 1 relies on new chain rules of quantum relative entropies. For classical probability distributions P_{XY} and Q_{XY} , the chain rule [42, Theorem 2.5.3] states that $D(P_{XY}||Q_{XY}) = D(P_X||Q_X) + \sum_x P_X(x)D(P_{Y|X=x}||Q_{Y|X=x})$, where D is the Kullback-Leibler divergence [43]. While no exact quantum analog exists, inequalities such as $D(P_{XY}||Q_{XY}) \leq D(P_X||Q_X) + \max_x D(P_{Y|X=x}||Q_{Y|X=x})$ have been extended to quantum settings for Umegaki [11] and Belavkin-Staszewski relative entropies [21]. Here, we establish a reverse inequality, providing the *first* quantum analog for $D(P_{XY}||Q_{XY}) \geq D(P_X||Q_X) + \min_x D(P_{Y|X=x}||Q_{Y|X=x})$.

The *measured relative entropy* is defined as [34, 44] $D_{\mathcal{M}}(\rho||\sigma) := \sup_{(\mathcal{X}, M)} D(P_{\rho, M}||P_{\sigma, M})$, where the supremum is over all finite sets \mathcal{X} and POVMs $M = \{M_x\}_{x \in \mathcal{X}}$ satisfying $M_x \geq 0$ and $\sum_{x \in \mathcal{X}} M_x = I$, and $P_{\rho, M}(x) := \text{Tr}[M_x \rho]$. The *measured Rényi divergence* is [45] $D_{\mathcal{M}, \alpha}(\rho||\sigma) := \sup_{(\mathcal{X}, M)} D_{\alpha}(P_{\rho, M}||P_{\sigma, M})$, where D_{α} is the classical Rényi divergence. The *sandwiched Rényi divergence* is [46, 47] $D_{\mathcal{S}, \alpha}(\rho||\sigma) := \frac{1}{\alpha-1} \log \text{Tr}[\sigma^{\frac{1-\alpha}{2\alpha}} \rho \sigma^{\frac{1-\alpha}{2\alpha}}]^{\alpha}$, if $\text{supp}(\rho) \subseteq \text{supp}(\sigma)$, and $+\infty$ otherwise. Then we have the chain rules as follows.

Lemma 2 (*Chain rules.*) *Let $\mathcal{N} \in \text{CPTP}(A : B)$, $\mathcal{M} \in \text{CP}(A : B)$, $\rho \in \mathcal{D}(RA)$, and $\sigma \in \mathcal{H}_+(RA)$. Let $\rho_{RB}^{\mathcal{N}} = \mathcal{N}(\rho_{RA})$ and $\sigma_{RB}^{\mathcal{M}} = \mathcal{M}(\sigma_{RA})$ be the channel output states. Then it holds, for any $\alpha \in (0, +\infty)$, that*

$$D_{\mathcal{M}, \alpha}(\rho_{RB}^{\mathcal{N}}||\sigma_{RB}^{\mathcal{M}}) \geq D_{\mathcal{M}, \alpha}(\rho_R||\sigma_R) + D_{\mathcal{M}, \alpha}^{\text{inf}}(\mathcal{N}||\mathcal{M}).$$

Moreover, it holds, for any $\alpha \in [1/2, \infty)$, that

$$D_{\mathcal{S}, \alpha}(\rho_{RB}^{\mathcal{N}}||\sigma_{RB}^{\mathcal{M}}) \geq D_{\mathcal{S}, \alpha}(\rho_R||\sigma_R) + D_{\mathcal{S}, \alpha}^{\text{inf}, \infty}(\mathcal{N}||\mathcal{M}).$$

When $\alpha = 1$, the chain rules represent the results for the measured and Umegaki relative entropies, respectively.

These chain rules can be seen as an enhancement of the data processing inequality under partial trace. Notably, they are also tight in the sense

that $\inf_{\rho, \sigma \in \mathcal{D}(RA)} [D_{s, \alpha}(\rho_{RB}^N \| \sigma_{RB}^M) - D_{s, \alpha}(\rho_R \| \sigma_R)] = D_{s, \alpha}^{\text{inf}, \infty}(\mathcal{N} \| \mathcal{M})$. That is, the amortized channel divergence coincides with the regularized divergence, serving as an analog to [11, Corollary 3] in the worst-case scenario. Detailed proofs of the chain rules and their tightness are provided in the Supplemental Materials.

Let $D_{H, \varepsilon}(\rho \| \sigma) := -\log\{\text{Tr}[\sigma M] : \text{Tr}[\rho(I - M)] \leq \varepsilon, 0 \leq M \leq I\}$ be the quantum hypothesis testing relative entropy. The proof for Theorem 1 then contains two parts. The converse part uses the chain rules to show that $\liminf_{n \rightarrow \infty} \frac{1}{n} D_{H, \varepsilon}(\mathcal{A}_n \| \mathcal{B}_n) \geq D^{\text{inf}, \infty}(\mathcal{N} \| \mathcal{M})$ while the achievable part shows that $\limsup_{n \rightarrow \infty} \frac{1}{n} D_{H, \varepsilon}(\mathcal{A}'_n \| \mathcal{B}'_n) \leq D^{\text{inf}, \infty}(\mathcal{N} \| \mathcal{M})$ by applying the generalized AEP in [41] to \mathcal{A}'_n and \mathcal{B}'_n .

1) *Proof of the converse part.* The first step is to lower bound $D_{H, \varepsilon}(\mathcal{A}_n \| \mathcal{B}_n)$ by applying the chain rule recursively. Denote the joint states before the n -th use of the channel by $\rho'_n := \text{Tr}_{R_n} \circ \mathcal{P}^n \circ \prod_{i=1}^{n-1} \mathcal{U} \circ \mathcal{P}^i$ and $\sigma'_n := \text{Tr}_{R_n} \circ \mathcal{Q}^n \circ \prod_{i=1}^{n-1} \mathcal{V} \circ \mathcal{Q}^i$. Denote $\rho_n := \rho[\{\mathcal{P}^i\}_{i=1}^n]$ and $\sigma_n := \sigma[\{\mathcal{Q}^i\}_{i=1}^n]$. Then we have that $\rho_n = \text{Tr}_{R_n E_n} \circ \mathcal{U}(\rho'_n) = \mathcal{N}(\rho'_n)$ and $\sigma_n = \text{Tr}_{R_n E_n} \circ \mathcal{V}(\sigma'_n) = \mathcal{M}(\sigma'_n)$. Note that for any \mathcal{P}^n and \mathcal{Q}^n we always have $\text{Tr}_{A_n R_n} \circ \mathcal{P}^n = \text{Tr}_{E_{n-1} R_{n-1}}$ and $\text{Tr}_{A_n R_n} \circ \mathcal{Q}^n = \text{Tr}_{E_{n-1} R_{n-1}}$. This gives the relations that $\text{Tr}_{A_n}(\rho'_n) = \rho_{n-1}$ and $\text{Tr}_{A_n}(\sigma'_n) = \sigma_{n-1}$. Applying the chain rule in Lemma 2 to ρ_n and σ_n , we have

$$D_{s, \alpha}(\rho_n \| \sigma_n) \geq D_{s, \alpha}(\rho_{n-1} \| \sigma_{n-1}) + D_{s, \alpha}^{\text{inf}, \infty}(\mathcal{N} \| \mathcal{M}).$$

Recursively applying this relation n times, we get

$$D_{s, \alpha}(\rho_n \| \sigma_n) \geq n D_{s, \alpha}^{\text{inf}, \infty}(\mathcal{N} \| \mathcal{M}).$$

As this holds for any operations \mathcal{P}^i and \mathcal{Q}^i , we get

$$D_{s, \alpha}(\mathcal{A}_n \| \mathcal{B}_n) \geq n D_{s, \alpha}^{\text{inf}, \infty}(\mathcal{N} \| \mathcal{M}).$$

Using the relation of hypothesis testing relative entropy and the Petz Rényi divergence in [48, Proposition 3] and the fact that Petz Rényi divergence is no smaller than the sandwiched Rényi divergence [49, Eq.(4.88)], we have for any $\alpha \in [1/2, 1)$ and $\varepsilon \in (0, 1)$, that

$$D_{H, \varepsilon}(\mathcal{A}_n \| \mathcal{B}_n) \geq D_{s, \alpha}(\mathcal{A}_n \| \mathcal{B}_n) + (\log(1/\varepsilon))\alpha/(\alpha - 1).$$

Combining the above relations, we have

$$D_{H, \varepsilon}(\mathcal{A}_n \| \mathcal{B}_n) \geq n D_{s, \alpha}^{\text{inf}, \infty}(\mathcal{N} \| \mathcal{M}) + (\log(1/\varepsilon))\alpha/(\alpha - 1).$$

Taking the limits on both sides, we get

$$\liminf_{n \rightarrow \infty} \frac{1}{n} D_{H, \varepsilon}(\mathcal{A}_n \| \mathcal{B}_n) \geq \sup_{\alpha \in [1/2, 1)} D_{s, \alpha}^{\text{inf}, \infty}(\mathcal{N} \| \mathcal{M}).$$

Finally, note that $\sup_{\alpha \in [1/2, 1)} D_{s, \alpha}^{\text{inf}, \infty}(\mathcal{N} \| \mathcal{M}) \geq \sup_{\alpha \in [1/2, 1)} D_{M, \alpha}^{\text{inf}, \infty}(\mathcal{N} \| \mathcal{M}) = D^{\text{inf}, \infty}(\mathcal{N} \| \mathcal{M})$, where the inequality follows as $D_{M, \alpha}(\rho \| \sigma) \leq D_{s, \alpha}(\rho \| \sigma)$ for

$\alpha \in [1/2, 1)$, and the equality is a consequence of [41, Lemma 27, 28], applied to the image sets of the channels. This concludes the proof of the converse part.

2) *Proof of the achievable part.* We check that the sets $\{\mathcal{A}'_n\}_n$ and $\{\mathcal{B}'_n\}_n$ meet all the structural assumptions to apply the generalized AEP in [41, Theorem 25]. First, the set of all density matrices \mathcal{D} is convex and compact, so \mathcal{A}'_n is also convex and compact. Since $\mathcal{N}^{\otimes n}$ and \mathcal{D} are permutation invariant, we know that \mathcal{A}'_n is also permutation invariant. For any $\mathcal{N}^{\otimes m}(\rho_m) \in \mathcal{A}'_m$ and $\mathcal{N}^{\otimes k}(\rho_k) \in \mathcal{A}'_k$, we have $\mathcal{N}^{\otimes m}(\rho_m) \otimes \mathcal{N}^{\otimes k}(\rho_k) = \mathcal{N}^{\otimes(m+k)}(\rho_m \otimes \rho_k) \in \mathcal{A}'_{m+k}$. This implies $\mathcal{A}'_m \otimes \mathcal{A}'_k \subseteq \mathcal{A}'_{m+k}$. The support function of \mathcal{A}'_n is given by $h_{\mathcal{A}'_n}(X_n) := \sup_{\rho_n \in \mathcal{D}} \text{Tr}[X_n \mathcal{N}^{\otimes n}(\rho_n)] = \sup_{\rho_n \in \mathcal{D}} \text{Tr}[(\mathcal{N}^{\otimes n})^\dagger(X_n) \rho_n] = \lambda_{\max}((\mathcal{N}^{\otimes n})^\dagger(X_n))$ where $\lambda_{\max}(\cdot)$ denotes the maximum eigenvalue. Therefore, for any $X_m \in \mathcal{H}_+$ and $X_k \in \mathcal{H}_+$, we have $h_{\mathcal{A}'_{m+k}}(X_m \otimes X_k) = h_{\mathcal{A}'_m}(X_m) h_{\mathcal{A}'_k}(X_k)$, by the multiplicativity of the maximum eigenvalue under tensor product. This proves that $\{\mathcal{A}'_n\}_{n \in \mathbb{N}}$ satisfy all the required assumptions, and the same argument works for $\{\mathcal{B}'_n\}_{n \in \mathbb{N}}$. This gives $\limsup_{n \rightarrow \infty} \frac{1}{n} D_{H, \varepsilon}(\mathcal{A}'_n \| \mathcal{B}'_n) = D^{\text{inf}, \infty}(\mathcal{N} \| \mathcal{M})$ from [41] and proves the achievable part.

Relative entropy accumulation.— The entropy accumulation theorem [50, 51] is a technique to find bounds on the operationally relevant uncertainty present in the outputs of a sequential process as a sum of the worst case uncertainties of each step. It has been widely used in quantum cryptography [36–38]. More specifically, a variant for the max-entropy H_{\max}^ε states that [51] for channels $\mathcal{N}_i \in \text{CPTP}(Y_{i-1} : Y_i S_i C_i)$, we have $H_{\max}^\varepsilon(S_1 \dots S_n | C_1 \dots C_n)_{\mathcal{N}_1 \circ \dots \circ \mathcal{N}_n}(\rho_{Y_0}) \leq \sum_{i=1}^n \sup_{\omega_{Y_{i-1}}} H(S_i | C_i)_{\mathcal{N}_i(\omega)} + O(\sqrt{n})$, where $H_{\max}^\varepsilon(S | C)_\rho \approx -\inf_{\sigma \in \mathcal{D}(C)} D_{H, \varepsilon}(\rho_{SC} \| I_S \otimes \sigma_C)$ [52] and $H(S | C)_\rho = -\inf_{\sigma \in \mathcal{D}(C)} D(\rho_{SC} \| I_S \otimes \sigma_C)$. This naturally raises the question of whether such a statement can be generalized to divergences between arbitrary sequential processes of channels, rather than being restricted only to entropies (which corresponds to choosing \mathcal{M} in Figure 1 to be a replacer channel). This was first asked as an open question in [51] for the max-relative entropy. By extending the adversarial channel discrimination framework in Figure 1 to allow different channels \mathcal{N}_i and \mathcal{M}_i at each round, we can unify the relative entropy accumulation within this broader framework. This new perspective enables us to establish a relative entropy accumulation theorem for $D_{H, \varepsilon}$ (a smoothed form of the min-relative entropy), giving an answer to the dual formulation of this open question. Specifically, we generalize the converse part of Theorem 1 in two ways: we allow the channels applied at different steps to vary, and we compute explicit finite-size bounds.

Theorem 3 (*Relative entropy accumulation theorem.*)
Let $\mathcal{N}_i \in \text{CPTP}(A_i : A_{i+1} B_i)$, $\mathcal{M}_i \in \text{CP}(A_i : A_{i+1} B_i)$

and $\rho, \sigma \in \mathcal{D}(A_1)$. Let $\varepsilon \in (0, 1)$, then it holds that

$$D_{H,\varepsilon} \left(\text{Tr}_{A_{n+1}} \circ \prod_{i=1}^n \mathcal{N}_i(\rho_{A_1}) \left\| \text{Tr}_{A_{n+1}} \circ \prod_{i=1}^n \mathcal{M}_i(\sigma_{A_1}) \right. \right) \geq \sum_{i=1}^n D^{\text{inf},\infty}(\text{Tr}_{A_{i+1}} \circ \mathcal{N}_i \parallel \text{Tr}_{A_{i+1}} \circ \mathcal{M}_i) - O(n^{2/3} \log n).$$

The proof makes important use of the chain rules in Lemma 2. Moreover, choosing the channels \mathcal{M}_i to be replacer channels, we recover a slightly weaker version of the H_{\max}^ε entropy accumulation statement previously mentioned. We leave it as an open question for future work whether this new proof technique can lead to better entropy accumulation theorems.

Discussion.— We introduced the adversarial quantum channel discrimination and established a direct analog of quantum Stein’s lemma in this new setting. Notably, the optimal error exponent can be efficiently computed despite its regularization, and the strong converse property holds in general. Moreover, the equivalence of adaptive and non-adaptive strategies by the adversary provides significant insight for the tester, as their optimal measurements can be performed as if playing against a much weaker non-adaptive adversary. We also derived a relative entropy accumulation theorem, extending the existing entropy accumulation theorem from entropy to divergence and providing a solution to the dual formulation of the open problem presented in [51]. On the technical side, we introduced the minimum output channel divergence and established its chain rules. This tool has recently been used in [53] for the security analysis of quantum cryptographic protocols. Given the fundamental importance of quantum channel discrimination in quantum information, we anticipate that the new framework and tools developed in this work will open new directions for future investigations.

Acknowledgments. K.F. is supported by the NSFC (grant No. 92470113 and 12404569), the Shenzhen Science and Technology Program (grant No. JCYJ20240813113519025). O.F. acknowledges support by the European Research Council (ERC Grant Algo-QIP, Agreement No. 851716), by the European Union’s Horizon research and innovation programme under the project VERIQ-TAS (Grant Agreement No 101017733) and the project “Quantum Security Networks Partnership” QSNP (Grant Agreement No 101114043), and by the Agence Nationale de la Recherche under the Plan France 2030 with the reference ANR-22-PETQ-0009.

* kunfang@cuhk.edu.cn

† h.fawzi@damtp.cam.ac.uk

‡ omar.fawzi@ens-lyon.fr

[1] A. Acín, *Physical Review Letters* **87**, 177901 (2001).

- [2] G. M. D’Ariano, *Physical Review Letters* **87** (2001).
- [3] G. Chiribella, G. M. D’Ariano, and P. Perinotti, *Physical Review Letters* **101**, 180501 (2008).
- [4] M. Hayashi, *IEEE Transactions on Information Theory* **55**, 3807 (2009).
- [5] A. W. Harrow, A. Hassidim, D. W. Leung, and J. Watrous, *Physical Review A* **81**, 032339 (2010).
- [6] M. M. Wilde, M. Berta, C. Hirche, and E. Kaur, *Letters in Mathematical Physics* **110**, 2277 (2020).
- [7] T. Cooney, M. Mosonyi, and M. M. Wilde, *Communications in Mathematical Physics* **344**, 797 (2016).
- [8] X. Wang and M. M. Wilde, *Physical Review Research* **1**, 033170 (2019).
- [9] S. Pirandola, R. Laurenza, C. Lupo, and J. L. Pereira, *npj Quantum Information* **5**, 50 (2019).
- [10] X. Wang and M. M. Wilde, *Physical Review Research* **1**, 033169 (2019).
- [11] K. Fang, O. Fawzi, R. Renner, and D. Sutter, *Physical Review Letters* **124**, 100501 (2020).
- [12] B. Bergh, N. Datta, R. Salzmänn, and M. M. Wilde, *IEEE Transactions on Information Theory* **70**, 2617 (2024).
- [13] M. Piani and J. Watrous, *Physical Review Letters* **102**, 250501 (2009).
- [14] R. Takagi, B. Regula, K. Bu, Z.-W. Liu, and G. Adesso, *Physical Review Letters* **122**, 140402 (2019).
- [15] J. Bae, D. Chruściński, and M. Piani, *Physical Review Letters* **122**, 140404 (2019).
- [16] P. Skrzypczyk and N. Linden, *Physical Review Letters* **122**, 140403 (2019).
- [17] L. Wang and R. Renner, *Physical Review Letters* **108**, 200501 (2012).
- [18] N. Datta, M. Mosonyi, M.-H. Hsieh, and F. G. S. L. Brandao, *IEEE Transactions on Information Theory* **59**, 8014 (2013).
- [19] X. Wang, K. Fang, and R. Duan, *IEEE Transactions on Information Theory* **65**, 2583 (2019).
- [20] X. Wang, K. Fang, and M. Tomamichel, *IEEE Transactions on Information Theory* **65**, 4609 (2019).
- [21] K. Fang and H. Fawzi, *Communications in Mathematical Physics* **384**, 1615 (2021).
- [22] K. Fang, G. Gour, and X. Wang, *arXiv preprint arXiv:2110.14842* (2021).
- [23] S. Pirandola, B. R. Bardhan, T. Gehring, C. Weedbrook, and S. Lloyd, *Nature Photonics* **12**, 724 (2018).
- [24] Q. Zhuang and S. Pirandola, *Physical Review Letters* **125**, 080505 (2020).
- [25] G. Spedalieri, L. Piersimoni, O. Laurino, S. L. Braunstein, and S. Pirandola, *Physical Review Research* **2**, 043260 (2020).
- [26] J. Pereira, *Quantum channel simulation and discrimination with applications to quantum communications*, Ph.D. thesis, University of York (2020).
- [27] C. H. Bennett and G. Brassard, in *International Conference on Computers, Systems & Signal Processing, Bangalore, India, Dec 9-12, 1984* (1984) pp. 175–179.
- [28] D. Mayers and A. Yao, in *Proceedings 39th Annual Symposium on Foundations of Computer Science (Cat. No. 98CB36280)* (IEEE, 1998) pp. 503–509.
- [29] S. Lu, L.-M. Duan, and D.-L. Deng, *Physical Review Research* **2**, 033212 (2020).
- [30] T. Vidick, J. Watrous, *et al.*, *Foundations and Trends® in Theoretical Computer Science* **11**, 1 (2016).

- [31] H. Zhu and M. Hayashi, *Physical Review Letters* **123**, 260504 (2019).
- [32] F. G. Brandão, A. W. Harrow, J. R. Lee, and Y. Peres, *IEEE Transactions on Information Theory* **66**, 5037 (2020).
- [33] In [32], the problem is formulated in terms of convex sets P, Q of probability distributions. For the quantum generalization (which is operationally well-motivated), we define the sets P and Q as the images of all input probability distributions of certain physical channels, denoted \mathcal{N} and \mathcal{M} . This means that the channel \mathcal{N} takes as input a description of some $p \in P$ and outputs a sample from p . If P is the convex hull of a finite set of probability distributions (p_1, \dots, p_k) , such a map \mathcal{N} can be constructed by letting $\mathcal{N}(|i\rangle\langle j|) = \delta_{i=j}p_i$. For more general convex sets, the channel \mathcal{N} would require an infinite-dimensional input, which is beyond the scope of this paper.
- [34] F. Hiai and D. Petz, *Communications in Mathematical Physics* **143**, 99 (1991).
- [35] H. Nagaoka and T. Ogawa, *IEEE Transactions on Information Theory* **46**, 2428 (2000).
- [36] R. Arnon-Friedman, F. Dupuis, O. Fawzi, R. Renner, and T. Vidick, *Nature Communications* **9**, 459 (2018).
- [37] Y. Liu, Q. Zhao, M.-H. Li, J.-Y. Guan, Y. Zhang, B. Bai, W. Zhang, W.-Z. Liu, C. Wu, X. Yuan, *et al.*, *Nature* **562**, 548 (2018).
- [38] I. George, J. Lin, T. van Himbeek, K. Fang, and N. Lütkenhaus, *arXiv preprint arXiv:2203.06554* (2022).
- [39] H. Umegaki, in *Kodai Mathematical Seminar Reports*, Vol. 14 (Department of Mathematics, Tokyo Institute of Technology, 1962) pp. 59–85.
- [40] K. Fang, H. Fawzi, and O. Fawzi, *arXiv preprint arXiv:2502.15659* (2025).
- [41] K. Fang, H. Fawzi, and O. Fawzi, *arXiv preprint arXiv:2411.04035* (2024).
- [42] T. M. Cover, *Elements of Information Theory* (John Wiley & Sons, 1999).
- [43] S. Kullback and R. A. Leibler, *The Annals of Mathematical Statistics* **22**, 79 (1951).
- [44] M. J. Donald, *Communications in Mathematical Physics* **105**, 13 (1986).
- [45] M. Berta, O. Fawzi, and M. Tomamichel, *Letters in Mathematical Physics* **107**, 2239 (2017).
- [46] M. Müller-Lennert, F. Dupuis, O. Szechr, S. Fehr, and M. Tomamichel, *Journal of Mathematical Physics* **54** (2013).
- [47] M. M. Wilde, A. Winter, and D. Yang, *Communications in Mathematical Physics* **331**, 593 (2014).
- [48] H. Qi, Q. Wang, and M. M. Wilde, *Journal of Physics A: Mathematical and Theoretical* **51**, 444002 (2018).
- [49] M. Tomamichel, *Quantum Information Processing with Finite Resources*, SpringerBriefs in Mathematical Physics, Vol. 5 (Springer International Publishing, Cham, 2016).
- [50] F. Dupuis, O. Fawzi, and R. Renner, *Communications in Mathematical Physics* **379**, 867 (2020).
- [51] T. Metger, O. Fawzi, D. Sutter, and R. Renner, in *2022 IEEE 63rd Annual Symposium on Foundations of Computer Science (FOCS)* (IEEE, 2022) pp. 844–850.
- [52] The exact definition is not in terms of $D_{H,\epsilon}$ but a closely related smoothed version of $D_{S,1/2}$. See Supplemental Materials for more details.
- [53] A. Arqand and E. Y.-Z. Tan, *arXiv preprint arXiv:2502.02563* (2025).
- [54] N. Datta, *IEEE Transactions on Information Theory* **55**, 2816 (2009).
- [55] R. Renner, *Security of QKD*, Ph.D. thesis, ETH, 2005, quant-ph/0512258 (2005).
- [56] D. Petz, *Reports on Mathematical Physics* **23**, 57 (1986).
- [57] J. Watrous, *The theory of quantum information* (Cambridge University Press, 2018).
- [58] F. Leditzky, E. Kaur, N. Datta, and M. M. Wilde, *Physical Review A* **97**, 012332 (2018).
- [59] M. Tomamichel, M. Berta, and M. Hayashi, *Journal of Mathematical Physics* **55** (2014).
- [60] F. Dupuis and O. Fawzi, *IEEE Transactions on Information Theory* **65**, 7596 (2019).
- [61] E. Carlen, *Entropy and the quantum* **529**, 73 (2010).
- [62] F. Dupuis, L. Kraemer, P. Faist, J. M. Renes, and R. Renner, in *XVIIth International Congress on Mathematical Physics* (World Scientific, 2014) pp. 134–153.
- [63] J. Watrous, *Theory of Computing* **5**, 217 (2009).

Supplemental Materials

In this Supplemental Material, we provide more detailed expositions, proofs and discussions of the results in the main text. We may reiterate some of the steps to ensure that the Supplemental Material are explicit and self-contained.

CONTENTS

Acknowledgments	5
References	5
Preliminaries	7
Notations and quantum divergences	7
Minimum output channel divergence	8
Chain rules	9
Proof of the chain rules	9
Tightness of the chain rules	10
Counter-example to a potential improvement of the chain rule	11
Proof of the adversarial quantum Stein's lemma	12
Relative entropy accumulation	13
Useful properties	18

PRELIMINARIES

Notations and quantum divergences

Let $\mathcal{D}(A)$ denote the set of all density operators on a finite-dimensional Hilbert space \mathcal{H}_A . Let $\mathcal{L}(A)$ represent the set of all linear operators. Let $\mathcal{H}(A)$, $\mathcal{H}_+(A)$ and $\mathcal{H}_{++}(A)$ be the set of all hermitian operators, positive semidefinite operators and positive definite operators on \mathcal{H}_A , respectively. The set of completely positive maps from $\mathcal{L}(A)$ to $\mathcal{L}(B)$ is denoted by $\text{CP}(A : B)$. A quantum channel $\mathcal{N}_{A \rightarrow B}$ is a linear map from $\mathcal{L}(A)$ to $\mathcal{L}(B)$ that is both completely positive and trace-preserving. The set of all such maps is denoted by $\text{CPTP}(A : B)$. A quantum divergence is a functional $\mathbb{D} : \mathcal{D} \times \mathcal{H}_+ \rightarrow \mathbb{R}$ that satisfies the data-processing inequality, characterizing the “distinguishability” or “distance” between two quantum states. There are a few quantum divergences used throughout this work.

Definition 1 (*Umegaki relative entropy [39].*) For any $\rho \in \mathcal{D}$ and $\sigma \in \mathcal{H}_+$, the Umegaki relative entropy is defined by

$$D(\rho \parallel \sigma) := \text{Tr}[\rho(\log \rho - \log \sigma)], \quad (2)$$

if $\text{supp}(\rho) \subseteq \text{supp}(\sigma)$ and $+\infty$ otherwise.

Definition 2 (*Sandwiched Rényi divergence [46, 47].*) Let $\alpha \in (0, 1) \cup (1, +\infty)$. For any $\rho \in \mathcal{D}$ and $\sigma \in \mathcal{H}_+$, the sandwiched Rényi divergence is defined by

$$D_{s,\alpha}(\rho \parallel \sigma) := \frac{1}{\alpha - 1} \log \text{Tr} \left[\sigma^{\frac{1-\alpha}{2\alpha}} \rho \sigma^{\frac{1-\alpha}{2\alpha}} \right]^\alpha, \quad (3)$$

if $\text{supp}(\rho) \subseteq \text{supp}(\sigma)$, and $+\infty$ otherwise.

When $\alpha \rightarrow 1$, $D_{s,\alpha}$ converge to the Umegaki relative entropy [46, 47],

$$\lim_{\alpha \rightarrow 1} D_{s,\alpha}(\rho \parallel \sigma) = D(\rho \parallel \sigma). \quad (4)$$

When $\alpha \rightarrow \infty$, the sandwiched Rényi divergence converges to the max-relative entropy [54, 55],

$$\lim_{\alpha \rightarrow \infty} D_{s,\alpha}(\rho \parallel \sigma) = D_{\max}(\rho \parallel \sigma) := \log \inf \{ t \in \mathbb{R} : \rho \leq t\sigma \}, \quad (5)$$

if $\text{supp}(\rho) \subseteq \text{supp}(\sigma)$ and $+\infty$ otherwise.

Definition 3 (Petz Rényi divergence [56].) *Let $\alpha \in (0, 1) \cup (1, +\infty)$. For any $\rho \in \mathcal{D}$ and $\sigma \in \mathcal{H}_+$, the Petz Rényi divergence is defined by*

$$D_{P,\alpha}(\rho \parallel \sigma) := \frac{1}{\alpha - 1} \log \text{Tr} [\rho^\alpha \sigma^{1-\alpha}], \quad (6)$$

if $\text{supp}(\rho) \subseteq \text{supp}(\sigma)$, and $+\infty$ otherwise.

Definition 4 (Hypothesis testing relative entropy.) *Let $\varepsilon \in [0, 1]$. For any $\rho \in \mathcal{D}$ and $\sigma \in \mathcal{H}_+$, the quantum hypothesis testing relative entropy is defined by $D_{H,\varepsilon}(\rho \parallel \sigma) := -\log \beta_\varepsilon(\rho \parallel \sigma)$ where*

$$\beta_\varepsilon(\rho \parallel \sigma) := \min_{0 \leq M \leq I} \{ \text{Tr}[\sigma M] : \text{Tr}[\rho(I - M)] \leq \varepsilon \}. \quad (7)$$

Definition 5 (Measured relative entropy [34, 44].) *For any $\rho \in \mathcal{D}$, $\sigma \in \mathcal{H}_+$, the measured relative entropy is defined by*

$$D_M(\rho \parallel \sigma) := \sup_{(\mathcal{X}, M)} D(P_{\rho, M} \parallel P_{\sigma, M}), \quad (8)$$

where D is the Kullback–Leibler divergence and the optimization is over finite sets \mathcal{X} and positive operator valued measures M on \mathcal{X} such that $M_x \geq 0$ and $\sum_{x \in \mathcal{X}} M_x = I$, $P_{\rho, M}$ is a measure on \mathcal{X} defined via the relation $P_{\rho, M}(x) = \text{Tr}[M_x \rho]$ for any $x \in \mathcal{X}$.

Definition 6 (Measured Rényi divergence [45].) *Let $\alpha \in (0, 1) \cup (1, \infty)$. For any $\rho \in \mathcal{D}$ and $\sigma \in \mathcal{H}_+$, the measured Rényi divergence is defined as*

$$D_{M,\alpha}(\rho \parallel \sigma) := \sup_{(\mathcal{X}, M)} D_\alpha(P_{\rho, M} \parallel P_{\sigma, M}), \quad (9)$$

where D_α is the classical Rényi divergence.

When $\alpha \rightarrow 1$, the measured Rényi divergence converges to the measured relative entropy.

Minimum output channel divergence

The quantum divergence between two quantum states can be naturally extended to two sets of quantum states.

Definition 7 (Quantum divergence between two sets of states.) *For any subsets $\mathcal{A} \subseteq \mathcal{D}$ and $\mathcal{B} \subseteq \mathcal{H}_+$, their divergence is defined as*

$$\mathbb{D}(\mathcal{A} \parallel \mathcal{B}) := \inf_{\rho \in \mathcal{A}, \sigma \in \mathcal{B}} \mathbb{D}(\rho \parallel \sigma), \quad (10)$$

which represents the minimum “distance” between the sets.

Definition 8 (Minimum output channel divergence.) *Let $\mathcal{N} \in \text{CPTP}(A : B)$ and $\mathcal{M} \in \text{CP}(A : B)$. The minimum output channel divergence is defined by*

$$\mathbb{D}^{\text{inf}}(\mathcal{N} \parallel \mathcal{M}) := \inf_{\rho, \sigma \in \mathcal{D}(A)} \mathbb{D}(\mathcal{N}(\rho) \parallel \mathcal{M}(\sigma)), \quad (11)$$

which conceptually captures the worst-case scenario where different test states are chosen to minimize the distinguishability between the given channels.

It is also useful to see this as a divergence between two sets of quantum states, expressed as

$$\mathbb{D}^{\text{inf}}(\mathcal{N}\|\mathcal{M}) = \mathbb{D}(\mathcal{N}(\mathcal{D})\|\mathcal{M}(\mathcal{D})), \quad (12)$$

where $\mathcal{L}(\mathcal{D}) := \{\mathcal{L}(\rho) : \rho \in \mathcal{D}\}$ denotes the image set of \mathcal{D} under the linear map \mathcal{L} . We also define the regularized channel divergence as

$$\mathbb{D}^{\text{inf},\infty}(\mathcal{N}\|\mathcal{M}) := \lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{D}^{\text{inf}}(\mathcal{N}^{\otimes n}\|\mathcal{M}^{\otimes n}), \quad (13)$$

which accounts for the asymptotic behavior of the channel divergence over multiple uses.

Remark 1 *As proved in the main text that the image sets satisfy all assumptions in [41, Assumption 24]. Note that the measured Rényi divergence coincides with the sandwiched Rényi divergence at $\alpha \in \{1/2, +\infty\}$. Thus, by applying the superadditivity in [41, Lemma 21] and the subadditivity in [41, Lemma 26], we conclude that the minimum output channel divergence is additive for these cases, that is,*

$$D_{s,1/2}^{\text{inf}}(\mathcal{N}_1 \otimes \mathcal{N}_2\|\mathcal{M}_1 \otimes \mathcal{M}_2) = D_{s,1/2}^{\text{inf}}(\mathcal{N}_1\|\mathcal{M}_1) + D_{s,1/2}^{\text{inf}}(\mathcal{N}_2\|\mathcal{M}_2) \quad (14)$$

$$D_{\text{max}}^{\text{inf}}(\mathcal{N}_1 \otimes \mathcal{N}_2\|\mathcal{M}_1 \otimes \mathcal{M}_2) = D_{\text{max}}^{\text{inf}}(\mathcal{N}_1\|\mathcal{M}_1) + D_{\text{max}}^{\text{inf}}(\mathcal{N}_2\|\mathcal{M}_2), \quad (15)$$

where the first equality recovers Watrous' result [57, Corollary 3.60].

CHAIN RULES

Proof of the chain rules

We have the following chain rules for measured Rényi divergences and sandwiched Rényi divergences, which can also be seen as an enhancement of the data processing inequality under partial trace. The proof of these chain rules requires the notion of (reverse) polar sets.

Definition 9 *Let $\mathcal{C} \subseteq \mathcal{H}$ be a convex set. Its polar set is defined by*

$$\mathcal{C}^\circ := \{X : \text{Tr}[XY] \leq 1, \forall Y \in \mathcal{C}\} = \{X : h_{\mathcal{C}}(X) \leq 1\} \quad (16)$$

where $h_{\mathcal{C}}$ is the support function of \mathcal{C} :

$$h_{\mathcal{C}}(\omega) := \sup_{\sigma \in \mathcal{C}} \text{Tr}[\omega\sigma]. \quad (17)$$

Let $\mathcal{C}_+^\circ := \mathcal{C}^\circ \cap \mathcal{H}_+$ and $\mathcal{C}_{++}^\circ := \mathcal{C}^\circ \cap \mathcal{H}_{++}$ be the intersections with positive semidefinite operators and positive definite operators, respectively.

Similarly, the reverse polar set of \mathcal{C} is defined as:

$$\mathcal{C}^* := \{X : \text{Tr}[XY] \geq 1, \forall Y \in \mathcal{C}\} = \{X : \underline{h}_{\mathcal{C}}(X) \geq 1\} \quad (18)$$

where $\underline{h}_{\mathcal{C}}$ is the reverse support function of \mathcal{C} :

$$\underline{h}_{\mathcal{C}}(\omega) := \inf_{\sigma \in \mathcal{C}} \text{Tr}[\omega\sigma]. \quad (19)$$

Lemma 4 (Chain rules.) *Let $\rho \in \mathcal{D}(RA)$, $\sigma \in \mathcal{H}_+(RA)$, $\mathcal{N} \in \text{CPTP}(A:B)$ and $\mathcal{M} \in \text{CP}(A:B)$. Let $\rho_{RB}^{\mathcal{N}} = \mathcal{N}(\rho_{RA})$ and $\sigma_{RB}^{\mathcal{M}} = \mathcal{M}(\sigma_{RA})$ be the channel output states. Then it holds, for any $\alpha \in (0, +\infty)$, that*

$$D_{\text{M},\alpha}(\rho_{RB}^{\mathcal{N}}\|\sigma_{RB}^{\mathcal{M}}) \geq D_{\text{M},\alpha}(\rho_R\|\sigma_R) + D_{\text{M},\alpha}^{\text{inf}}(\mathcal{N}\|\mathcal{M}). \quad (20)$$

Moreover, it holds, for any $\alpha \in [1/2, \infty)$, that

$$D_{\text{s},\alpha}(\rho_{RB}^{\mathcal{N}}\|\sigma_{RB}^{\mathcal{M}}) \geq D_{\text{s},\alpha}(\rho_R\|\sigma_R) + D_{\text{s},\alpha}^{\text{inf},\infty}(\mathcal{N}\|\mathcal{M}). \quad (21)$$

When $\alpha = 1$, the chain rules represent the results for the measured and Umegaki relative entropies, respectively.

Proof. The proof utilizes the superadditivity of the divergence between two sets of quantum states, as established in [41, Lemma 21, 23]. To this end, we consider the following sets:

$$\mathcal{A}_1 = \{\rho_R\}, \quad \mathcal{A}_2 = \mathcal{N}(\mathcal{D}), \quad \mathcal{A}_3 = \{\rho_{RB}^{\mathcal{N}}\}, \quad (22)$$

$$\mathcal{B}_1 = \{\sigma_R\}, \quad \mathcal{B}_2 = \mathcal{M}(\mathcal{D}), \quad \mathcal{B}_3 = \{\sigma_{RB}^{\mathcal{M}}\}, \quad (23)$$

and verify that they meet the required assumptions. We do this for $\{\mathcal{A}_1, \mathcal{A}_2, \mathcal{A}_3\}$ and the same argument works for $\{\mathcal{B}_1, \mathcal{B}_2, \mathcal{B}_3\}$. For any $Y_B \in (\mathcal{A}_2)_+^\circ$, we have $\text{Tr}[Y_B \mathcal{N}(\rho)] \leq 1$ for any $\rho \in \mathcal{D}(A)$. This implies that $\mathcal{N}^\dagger(Y_B) \leq I_A$, with \mathcal{N}^\dagger being the adjoint map of \mathcal{N} . Therefore, for any $X_R \in (\mathcal{A}_1)_+^\circ$ and $Y_B \in (\mathcal{A}_2)_+^\circ$, we have the following equations,

$$\text{Tr}[(X_R \otimes Y_B) \mathcal{N}_{A \rightarrow B}(\rho_{RA})] = \text{Tr}[(X_R \otimes \mathcal{N}^\dagger(Y_B)) \rho_{RA}] \quad (24)$$

$$\leq \text{Tr}[(X_R \otimes I_A)(\rho_{RA})] \quad (25)$$

$$= \text{Tr}[X_R \rho_R] \quad (26)$$

$$\leq 1. \quad (27)$$

This implies that $X_R \otimes Y_B \in (\mathcal{A}_3)_+^\circ$ and therefore $(\mathcal{A}_1)_+^\circ \otimes (\mathcal{A}_2)_+^\circ \subseteq (\mathcal{A}_3)_+^\circ$. Applying the superadditivity in [41, Lemma 21, 23], we have the asserted result in Eq. (20) for $\alpha \in (0, 1]$. The proof for $\alpha \in (1, +\infty)$ follows in the same way. For $\alpha \in (1, \infty)$, we work with the reverse polar sets instead: if $Y_B \in (\mathcal{A}_2)_+^*$, then $\mathcal{N}^\dagger(Y_B) \geq I_A$ and as a result, if $X_R \in (\mathcal{A}_1)_+^*$, we have $X_R \otimes Y_B \in (\mathcal{A}_3)_+^*$ and we can similarly apply [41, Lemma 21]. The result in Eq. (21) is a direct consequence of Eq. (20). More specifically, we have that for any $\alpha \in [1/2, \infty)$,

$$D_{s,\alpha}(\rho_{RB}^{\mathcal{N}} \|\sigma_{RB}^{\mathcal{M}}) = \lim_{n \rightarrow \infty} \frac{1}{n} D_{M,\alpha}((\rho_{RB}^{\mathcal{N}})^{\otimes n} \| (\sigma_{RB}^{\mathcal{M}})^{\otimes n}) \quad (28)$$

$$\geq \lim_{n \rightarrow \infty} \frac{1}{n} D_{M,\alpha}((\rho_R)^{\otimes n} \| (\sigma_B)^{\otimes n}) + \lim_{n \rightarrow \infty} \frac{1}{n} D_{M,\alpha}^{\text{inf}}(\mathcal{N}^{\otimes n} \|\mathcal{M}^{\otimes n}) \quad (29)$$

$$= D_{s,\alpha}(\rho_R \|\sigma_R) + D_{s,\alpha}^{\text{inf},\infty}(\mathcal{N} \|\mathcal{M}), \quad (30)$$

where the first line follows from [41, Lemma 16, 17], the second line follows from Eq. (20) and the last line follows from [41, Lemma 28]. \square

Tightness of the chain rules

In the following, we introduce the notion of the amortized minimum output channel divergence and show that it coincides with the regularized divergence, being an analog result for the best-case channel divergence [11]. This, in turn, demonstrates the tightness of our chain rule properties.

Similar to the amortized channel divergence used in the existing literature [6], we can define the minimum output version as follows. Let \mathbb{D} be a quantum divergence between states. Let $\mathcal{N} \in \text{CPTP}(A:B)$ and $\mathcal{M} \in \text{CP}(A:B)$. Then the amortized minimum output channel divergence is defined by

$$\mathbb{D}^{\text{inf,amo}}(\mathcal{N} \|\mathcal{M}) := \inf_{\substack{\rho \in \mathcal{D}(RA) \\ \sigma \in \mathcal{D}(RA)}} \mathbb{D}(\mathcal{N}_{A \rightarrow B}(\rho_{RA}) \|\mathcal{M}_{A \rightarrow B}(\sigma_{RA})) - \mathbb{D}(\rho_R \|\sigma_R). \quad (31)$$

Lemma 5 *Let $\alpha \in [1/2, \infty)$. For any $\mathcal{N} \in \text{CPTP}(A:B)$, $\mathcal{M} \in \text{CP}(A:B)$, it holds that*

$$D_{s,\alpha}^{\text{inf,amo}}(\mathcal{N} \|\mathcal{M}) = D_{s,\alpha}^{\text{inf},\infty}(\mathcal{N} \|\mathcal{M}). \quad (32)$$

Equivalently, for any $\mathcal{N} \in \text{CPTP}(A:B)$, $\mathcal{M} \in \text{CP}(A:B)$ and any $\varepsilon \in (0, 1)$, there exists $\rho, \sigma \in \mathcal{D}(RA)$, such that

$$D_{s,\alpha}^{\text{inf},\infty}(\mathcal{N} \|\mathcal{M}) \leq D_{s,\alpha}(\mathcal{N}_{A \rightarrow B}(\rho_{RA}) \|\mathcal{M}_{A \rightarrow B}(\sigma_{RA})) - D_{s,\alpha}(\rho_R \|\sigma_R) \leq D_{s,\alpha}^{\text{inf},\infty}(\mathcal{N} \|\mathcal{M}) + \varepsilon. \quad (33)$$

Proof. We prove the result for the quantum relative entropy ($\alpha = 1$) and the same argument works for the sandwiched Rényi divergence of order $\alpha \neq 1$ as well. Note that the chain rule property in Lemma 4 is equivalent to $D^{\text{inf,amo}}(\mathcal{N} \|\mathcal{M}) \geq D^{\text{inf},\infty}(\mathcal{N} \|\mathcal{M})$. Now we prove the reverse direction. For this, we will first show the superadditivity of the amortized divergence under tensor product. That is,

$$D^{\text{inf,amo}}(\mathcal{N}_1 \otimes \mathcal{N}_2 \|\mathcal{M}_1 \otimes \mathcal{M}_2) \geq D^{\text{inf,amo}}(\mathcal{N}_1 \|\mathcal{M}_1) + D^{\text{inf,amo}}(\mathcal{N}_2 \|\mathcal{M}_2), \quad (34)$$

for any quantum channels $\mathcal{N}_1 \in \text{CPTP}(A_1 : B_1)$, $\mathcal{N}_2 \in \text{CPTP}(A_2 : B_2)$, CP maps $\mathcal{M}_1 \in \text{CP}(A_1 : B_1)$, $\mathcal{M}_2 \in \text{CP}(A_2 : B_2)$. To see this, let $(\rho_{RA_1A_2}, \sigma_{RA_1A_2})$ be any feasible solution to the optimization on the left-hand side of Eq. (34) and denote its corresponding objective value by

$$\delta_{12} := D(\mathcal{N}_1 \otimes \mathcal{N}_2(\rho_{RA_1A_2}) \| \mathcal{M}_1 \otimes \mathcal{M}_2(\sigma_{RA_1A_2})) - D(\rho_R \| \sigma_R). \quad (35)$$

Let $\omega_{RA_2B_1} = \mathcal{N}_1(\rho_{RA_1A_2})$ and $\gamma_{RA_2B_1} = \mathcal{M}_1(\sigma_{RA_1A_2})$. We can check that $(\rho_{RA_1}, \sigma_{RA_1})$ and $(\omega_{RA_2B_1}, \gamma_{RA_2B_1})$ are feasible solutions to the amortized divergence on the right-hand side of Eq. (34), respectively, with the corresponding objective values by

$$\delta_1 := D(\mathcal{N}_1(\rho_{RA_1}) \| \mathcal{M}_1(\sigma_{RA_1})) - D(\rho_R \| \sigma_R) \geq D^{\text{inf,amo}}(\mathcal{N}_1 \| \mathcal{M}_1) \quad (36)$$

$$\delta_2 := D(\mathcal{N}_2(\omega_{RA_2B_1}) \| \mathcal{M}_2(\gamma_{RA_2B_1})) - D(\omega_{RB_1} \| \gamma_{RB_1}) \geq D^{\text{inf,amo}}(\mathcal{N}_2 \| \mathcal{M}_2). \quad (37)$$

Noting that $\omega_{RB_1} = \mathcal{N}_1(\rho_{RA_1})$ and $\gamma_{RB_1} = \mathcal{M}_1(\sigma_{RA_1})$, we have $\delta_{12} = \delta_1 + \delta_2$. This implies

$$\delta_{12} \geq D^{\text{inf,amo}}(\mathcal{N}_1 \| \mathcal{M}_1) + D^{\text{inf,amo}}(\mathcal{N}_2 \| \mathcal{M}_2). \quad (38)$$

As this holds for any feasible solution $(\rho_{RA_1A_2}, \sigma_{RA_1A_2})$, we have the asserted result in Eq. (34).

By trivializing the reference system in the amortized divergence, we get $D^{\text{inf,amo}}(\mathcal{N} \| \mathcal{M}) \leq D^{\text{inf}}(\mathcal{N} \| \mathcal{M})$. Then we have

$$D^{\text{inf,amo}}(\mathcal{N} \| \mathcal{M}) \leq \frac{1}{n} D^{\text{inf,amo}}(\mathcal{N}^{\otimes n} \| \mathcal{M}^{\otimes n}) \leq \frac{1}{n} D^{\text{inf}}(\mathcal{N}^{\otimes n} \| \mathcal{M}^{\otimes n}), \quad (39)$$

where the first inequality follows from Eq. (34). As the above holds for any n , we can take $n \rightarrow \infty$ on the right-hand side and conclude that $D^{\text{inf,amo}}(\mathcal{N} \| \mathcal{M}) \leq D^{\text{inf},\infty}(\mathcal{N} \| \mathcal{M})$. This completes the proof. \square

Counter-example to a potential improvement of the chain rule

The quantum channel divergence studied in most existing literatures [58] use the same test states for both channels. So it may be expected that we can enhance the chain rules by using the same test states as well. However, we show here that this is not possible by giving a counter-example. That is, the chain rule cannot be enhanced to

$$D_{\text{M}}(\mathcal{N}_{A \rightarrow B}(\rho_{RA}) \| \mathcal{M}_{A \rightarrow B}(\sigma_{RA})) \geq D_{\text{M}}(\rho_R \| \sigma_R) + D_{\text{M}}^{\text{inf}'}(\mathcal{N} \| \mathcal{M}) \quad (40)$$

where the channel divergence takes the same input state

$$D_{\text{M}}^{\text{inf}'}(\mathcal{N} \| \mathcal{M}) := \inf_{\rho \in \mathcal{D}(A)} D_{\text{M}}(\mathcal{N}_{A \rightarrow B}(\rho_A) \| \mathcal{M}_{A \rightarrow B}(\rho_A)). \quad (41)$$

To see this, consider the generalized amplitude damping (GAD) channel, which is defined as

$$\mathcal{A}_{\gamma, N}(\rho) = \sum_{i=1}^4 A_i \rho A_i^\dagger, \quad \gamma, N \in [0, 1], \quad (42)$$

with the Kraus operator

$$A_1 = \sqrt{1-N}(|0\rangle\langle 0| + \sqrt{1-\gamma}|1\rangle\langle 1|), \quad A_2 = \sqrt{\gamma(1-N)}|0\rangle\langle 1|, \quad (43)$$

$$A_3 = \sqrt{N}(\sqrt{1-\gamma}|0\rangle\langle 0| + |1\rangle\langle 1|), \quad A_4 = \sqrt{\gamma N}|1\rangle\langle 0|. \quad (44)$$

Using convex optimization, we can numerically evaluate each terms $D_{\text{M}}(\mathcal{N}_{A \rightarrow B}(\rho_{RA}) \| \mathcal{M}_{A \rightarrow B}(\sigma_{RA}))$, $D_{\text{M}}(\rho_R \| \sigma_R)$ and $D_{\text{M}}^{\text{inf}'}(\mathcal{N} \| \mathcal{M})$. Then in Figure 2(a), we show that the channel divergence $D_{\text{M}}^{\text{inf}'}$ is subadditive under tensor product of channels. That is, it does not inherit the properties of the state divergence, making it not a suitable channel extension. Moreover, in Figure 2(b), we show that the chain rule property in Eq. (40) does not hold, as there are cases such that $y < x_2$ in the plot.




FIG. 2. (a) Subadditivity for the channel divergence $D^{\text{M,inf}'}$ where $\mathcal{A}_{0.5,0}$ and $\mathcal{A}_{p,0.9}$ are the GAD channels and p ranges from 0 to 1; (b) Random test for the chain rule property, where the quantum channels are chosen as $\mathcal{A}_{0.5,0}$ and $\mathcal{A}_{0.5,0.9}$, the quantum states are 500 randomly generated quantum states with real entries, $x_1 = D_{\text{M}}(\rho_R || \sigma_R) + D_{\text{M}}^{\text{inf}'}(\mathcal{A}_{0.5,0} || \mathcal{A}_{0.5,0.9})$ and $x_2 = D_{\text{M}}(\rho_R || \sigma_R) + D_{\text{M}}^{\text{inf}'}(\mathcal{A}_{0.5,0} || \mathcal{A}_{0.5,0.9})$.

PROOF OF THE ADVERSARIAL QUANTUM STEIN'S LEMMA

Since non-adaptive strategies are a specific type of adaptive strategy, we have the inclusions that $\mathcal{A}'_n \subseteq \mathcal{A}_n$ and $\mathcal{B}'_n \subseteq \mathcal{B}_n$. This gives the relations for the type-I and type-II errors by $\alpha(\mathcal{A}'_n, M_n) \leq \alpha(\mathcal{A}_n, M_n)$ and $\beta(\mathcal{B}'_n, M_n) \leq \beta(\mathcal{B}_n, M_n)$. So we have the general relation that $\beta'_{n,\varepsilon}(\mathcal{N} || \mathcal{M}) := \inf\{\beta(\mathcal{B}'_n, M_n) : 0 \leq M_n \leq I, \alpha(\mathcal{A}'_n, M_n) \leq \varepsilon\} \leq \beta_{n,\varepsilon}(\mathcal{N} || \mathcal{M})$ where the inequality holds because the right-hand side represents the infimum of a larger objective value over a smaller feasible set. We can also rewrite the optimal type-II errors as the hypothesis testing relative entropy between two sets,

$$-\log \beta_{n,\varepsilon}(\mathcal{N} || \mathcal{M}) = D_{\text{H},\varepsilon}(\mathcal{A}_n || \mathcal{B}_n), \quad (45)$$

$$-\log \beta'_{n,\varepsilon}(\mathcal{N} || \mathcal{M}) = D_{\text{H},\varepsilon}(\mathcal{A}'_n || \mathcal{B}'_n), \quad (46)$$

where $D_{\text{H},\varepsilon}(\rho || \sigma) := -\log\{\text{Tr}[\rho M] : \text{Tr}[\rho(I - M)] \leq \varepsilon, 0 \leq M \leq I\}$. This can be done by showing that $\mathcal{A}_n(\mathcal{A}'_n)$ and $\mathcal{B}_n(\mathcal{B}'_n)$ are convex sets (see Lemma 6 below) and applying [41, Lemma 31]. After this, the proof for the adversarial quantum Stein's lemma then contains two parts, as shown in the main text. The converse part makes use of the chain rule property to show that

$$\liminf_{n \rightarrow \infty} \frac{1}{n} D_{\text{H},\varepsilon}(\mathcal{A}_n || \mathcal{B}_n) \geq D^{\text{inf},\infty}(\mathcal{N} || \mathcal{M}), \quad (47)$$

while the achievable part aims to show that

$$\limsup_{n \rightarrow \infty} \frac{1}{n} D_{\text{H},\varepsilon}(\mathcal{A}'_n || \mathcal{B}'_n) \leq D^{\text{inf},\infty}(\mathcal{N} || \mathcal{M}), \quad (48)$$

by applying the generalized quantum asymptotic equipartition (AEP) property in [41, Theorem 25] to \mathcal{A}'_n and \mathcal{B}'_n .

Lemma 6 (Convexity.) *The sets $\{\mathcal{A}_n\}_{n \in \mathbb{N}}$ and $\{\mathcal{B}_n\}_{n \in \mathbb{N}}$ defined in the main text are convex.*

Proof. We prove the assumptions for $\{\mathcal{A}_n\}_{n \in \mathbb{N}}$ and the same reasoning works for $\{\mathcal{B}_n\}_{n \in \mathbb{N}}$ as well. Let $\{\mathcal{P}^i\}_{i=1}^n$ with systems R_1, \dots, R_n and $\{\bar{\mathcal{P}}^i\}_{i=1}^n$ with systems R_1, \dots, R_n be two strategies and $\lambda \in [0, 1]$. Note that we may assume both strategies have the same systems R_i as we can always increase the dimension of the systems R_i by extending the action of the channel in an arbitrary way without affecting the output. Let us now define another strategy $\{\bar{\mathcal{P}}^i\}_{i=1}^n$ as follows. Let $\bar{R}_i = R_i \otimes C$ for all $i = 1, \dots, n$ where C is a two-dimensional system. Then define

$$\bar{\mathcal{P}}^1(\cdot) = \lambda \mathcal{P}^1(\cdot) \otimes |0\rangle\langle 0|_C + (1 - \lambda) \bar{\mathcal{P}}^1(\cdot) \otimes |1\rangle\langle 1|_C \quad (49)$$

and for $i \geq 2$, define

$$\bar{\mathcal{P}}^i(X) = (\mathcal{P}^i \circ \mathcal{C}_0(X)) \otimes |0\rangle\langle 0|_C + (\bar{\mathcal{P}}^i \circ \mathcal{C}_1(X)) \otimes |1\rangle\langle 1|_C \quad (50)$$

where $\mathcal{C}_0(X) = \langle 0|_C X |0\rangle_C$ and $\mathcal{C}_1(X) = \langle 1|_C X |1\rangle_C$. Since $\mathcal{C}_0, \mathcal{C}_1, \mathcal{P}^i, \bar{\mathcal{P}}^i$ are all CP maps, we know that $\bar{\mathcal{P}}^i$ is also a CP map. Moreover, if $\mathcal{P}^i, \bar{\mathcal{P}}^i$ are trace-preserving, then $\bar{\mathcal{P}}^i$ is also trace-preserving. It is easy to check the following relations,

$$\mathcal{C}_0 \circ \bar{\mathcal{P}}^1 = \lambda \mathcal{P}^1 \quad \text{and} \quad \mathcal{C}_1 \circ \bar{\mathcal{P}}^1 = (1 - \lambda) \bar{\mathcal{P}}^1, \quad (51)$$

$$\mathcal{C}_0 \circ \bar{\mathcal{P}}^i = \mathcal{P}^i \circ \mathcal{C}_0 \quad \text{and} \quad \mathcal{C}_1 \circ \bar{\mathcal{P}}^i = \bar{\mathcal{P}}^i \circ \mathcal{C}_1, \quad \forall i \geq 2. \quad (52)$$

Noting that Tr_C commutes with \mathcal{U} as they are acting on different systems, we have

$$\text{Tr}_C \circ \prod_{i=1}^n \mathcal{U} \circ \bar{\mathcal{P}}^i = \mathcal{U} \circ \mathcal{P}^n \circ \mathcal{C}_0 \circ \prod_{i=1}^{n-1} \mathcal{U} \circ \bar{\mathcal{P}}^i + \mathcal{U} \circ \bar{\mathcal{P}}^n \circ \mathcal{C}_1 \circ \prod_{i=1}^{n-1} \mathcal{U} \circ \bar{\mathcal{P}}^i. \quad (53)$$

Also noting that \mathcal{C}_0 and \mathcal{C}_1 both commute with \mathcal{U} as they are acting on different systems and using the relations in Eqs. (51) and (52), we have

$$\mathcal{U} \circ \mathcal{P}^n \circ \mathcal{C}_0 \circ \prod_{i=1}^{n-1} \mathcal{U} \circ \bar{\mathcal{P}}^i = \lambda \mathcal{U} \circ \mathcal{P}^n \circ \prod_{i=1}^{n-1} \mathcal{U} \circ \mathcal{P}^i = \lambda \prod_{i=1}^n \mathcal{U} \circ \mathcal{P}^i, \quad (54)$$

$$\mathcal{U} \circ \bar{\mathcal{P}}^n \circ \mathcal{C}_1 \circ \prod_{i=1}^{n-1} \mathcal{U} \circ \bar{\mathcal{P}}^i = (1 - \lambda) \mathcal{U} \circ \mathcal{P}^n \circ \prod_{i=1}^{n-1} \mathcal{U} \circ \bar{\mathcal{P}}^i = (1 - \lambda) \prod_{i=1}^n \mathcal{U} \circ \bar{\mathcal{P}}^i. \quad (55)$$

Taking these into Eq. (53), we have

$$\text{Tr}_{\bar{R}_n E_n} \circ \prod_{i=1}^n \mathcal{U} \circ \bar{\mathcal{P}}^i = \lambda \text{Tr}_{R_n E_n} \circ \prod_{i=1}^n \mathcal{U} \circ \mathcal{P}^i + (1 - \lambda) \text{Tr}_{R_n E_n} \circ \prod_{i=1}^n \mathcal{U} \circ \bar{\mathcal{P}}^i \quad (56)$$

This shows that any mixture of the reduced states on $B_1 \cdots B_n$ by the strategies $\{\mathcal{P}^i\}_{i=1}^n$ and $\{\bar{\mathcal{P}}^i\}_{i=1}^n$ is also given by the reduced state of another strategy $\{\bar{\mathcal{P}}^i\}_{i=1}^n$ which proves the convexity of \mathcal{A}_n . \square

RELATIVE ENTROPY ACCUMULATION

The entropy accumulation theorem [50, 51] is a technique to find bounds on the operationally relevant uncertainty (entropy) present in the outputs of a sequential process as a sum of the worst case uncertainties (entropies) of each step. It has been widely used in quantum cryptography [36–38]. This naturally raises the question of whether such a statement can be generalized to divergences between arbitrary sequential processes of channels, rather than being restricted only to entropies. This was first asked as an open question in [51] for the max-relative entropy.

More specifically, the operational setting for relative entropy accumulation is depicted in Figure 3. Consider two states ρ_{A_1} and σ_{A_1} and quantum channels $\mathcal{N}_i \in \text{CPTP}(A_i : A_{i+1} B_i)$ and $\mathcal{M}_i \in \text{CP}(A_i : A_{i+1} B_i)$ that are applied sequentially from $i = 1$ to $i = n$ and generating the systems B_i . The systems A_i should be seen as an internal memory system that we do not control. The key question in the relative entropy accumulation asks: *Can we bound the operationally relevant divergence between the obtained states as the sum of the contributions of each step?*




FIG. 3. Illustration of the setting of the relative entropy accumulation.

Similar to the adversarial channel discrimination in the main text, we denote the Stinespring dilations of \mathcal{N}_i and \mathcal{M}_i as U_i and V_i , respectively, with the environmental system denoting as T_i . Denote also the corresponding channel as $\mathcal{U}_i(\cdot) = U_i(\cdot)U_i^\dagger$ and $\mathcal{V}_i(\cdot) = V_i(\cdot)V_i^\dagger$. Note that U_i is an isometry because \mathcal{N}_i is trace-preserving, but this is not necessarily the case for V_i . Then the setting in Figure 3 is equivalent to the diagram in Figure 4.




FIG. 4. Illustration of the setting of the relative entropy accumulation with Stinespring dilation.

By extending the adversarial channel discrimination framework to allow different channels \mathcal{N}_i and \mathcal{M}_i at each round, we can unify the relative entropy accumulation within this broader framework. More specifically, using a similar notation in the main text, we denote the Stinespring dilation of \mathcal{N}_i and \mathcal{M}_i as U_i and V_i , respectively. Denote also the corresponding channel as $\mathcal{U}_i(\cdot) = U_i(\cdot)U_i^\dagger$ and $\mathcal{V}_i(\cdot) = V_i(\cdot)V_i^\dagger$. An illustrative figure is given in Figure 5. Then, taking a particular choice,

$$\dim R_i = 1, \quad (57)$$

$$E_i = A_{i+1}T_i \quad (58)$$

$$\mathcal{P}^1 = \rho_{A_1}, \quad (59)$$

$$\mathcal{Q}^1 = \sigma_{A_1}, \quad (60)$$

$$\mathcal{P}^i = \mathcal{Q}^i = \text{Tr}_{T_{i-1}}, \quad \forall i \geq 2, \quad (61)$$

the adversarial discrimination framework in Figure 5 reduces to the relative entropy accumulation in Figure 4.




FIG. 5. Illustration of an extended setting of adversarial quantum channel discrimination.

This new perspective enables us to establish a relative entropy accumulation theorem for $D_{H,\varepsilon}$ (a smoothed form of the min-relative entropy), giving an answer to the dual formulation of the open question in [51]. Specifically, we generalize the converse part of the adversarial quantum Stein's lemma in two ways: we allow the channels applied at different steps to vary, and we compute explicit finite-size bounds.

We state the relative entropy accumulation theorem in this more general setting depicted in Figure 5. From the above discussion, it is clear that the version stated in the main text is a special case.

Theorem 7 (*Relative entropy accumulation.*) Consider two sequences of maps $\mathcal{U}_i \in \text{CPTP}(A_i : B_i E_i)$ and $\mathcal{V}_i \in \text{CP}(A_i : B_i E_i)$ for $i \in \{1, \dots, n\}$ and let $\mathcal{P}^i, \mathcal{Q}^i \in \text{CPTP}(E_{i-1} R_{i-1}, A_i R_i)$ as illustrated in Figure 5. Let $\varepsilon \in (0, 1)$ and assume for all $i \in \{1, \dots, n\}$ and $m \geq 2$,

$$\forall \alpha \in [1/2, 1] : D_{\text{P}, 3/2}(\rho_m^{(\alpha)} \| \sigma_m^{(\alpha)}) \leq \frac{C}{4} m \quad \text{and} \quad \forall \sigma \in \mathcal{D}(A_i) : \log \text{Tr}(\mathcal{V}_i(\sigma)) \leq \frac{C}{4}, \quad (62)$$

where $\rho_m^{(\alpha)}, \sigma_m^{(\alpha)}$ are outputs of the channels $(\text{Tr}_{E_i} \circ \mathcal{U}_i)^{\otimes m}, (\text{Tr}_{E_i} \circ \mathcal{V}_i)^{\otimes m}$ (respectively) and achieve the minimum $D_{\text{P}, \alpha}(\rho_m^{(\alpha)} \| \sigma_m^{(\alpha)}) = D_{\text{P}, \alpha}^{\text{inf}}((\text{Tr}_{E_i} \circ \mathcal{U}_i)^{\otimes m} \| (\text{Tr}_{E_i} \circ \mathcal{V}_i)^{\otimes m})$. Then letting $\rho_n = \text{Tr}_{R_n E_n} \prod_{i=1}^n \mathcal{U}_i \circ \mathcal{P}^i$ and $\sigma_n = \text{Tr}_{R_n E_n} \prod_{i=1}^n \mathcal{V}_i \circ \mathcal{Q}^i$, we have

$$D_{\text{H}, \varepsilon} \left(\rho_n \left\| \sigma_n \right. \right) \geq \sum_{i=1}^n D^{\text{inf}, \infty}(\text{Tr}_{E_i} \circ \mathcal{U}_i \| \text{Tr}_{E_i} \circ \mathcal{V}_i) - C' n^{2/3} \log n \log^{1/3} \frac{1}{\varepsilon}, \quad (63)$$

where C' is a constant that only depends on $d = \max_i \dim B_i$ and C .

Proof. The proof below is basically the same as the converse part of the adversarial quantum Stein's lemma in the main text. We start as usual by bounding the hypothesis testing relative entropy with a Rényi divergence of order $\alpha \in (0, 1)$:

$$D_{\text{H}, \varepsilon} \left(\rho_n \left\| \sigma_n \right. \right) \geq D_{\text{S}, \alpha} \left(\rho_n \left\| \sigma_n \right. \right) + \frac{\alpha}{\alpha - 1} \log(1/\varepsilon). \quad (64)$$

We can now introduce $\rho'_n = \text{Tr}_{R_n} \circ \mathcal{P}^n \circ \prod_{i=1}^{n-1} \mathcal{U}_i \circ \mathcal{P}^i$ and $\sigma'_n = \text{Tr}_{R_n} \circ \mathcal{Q}^n \circ \prod_{i=1}^{n-1} \mathcal{V}_i \circ \mathcal{Q}^i$ the joint states before the n -th use the channel $\mathcal{U}_i/\mathcal{V}_i$. We have $\rho_n = (\text{Tr}_{E_n} \circ \mathcal{U}_n)(\rho'_n)$. We use the chain rule for sandwiched relative entropy to bound

$$D_{\text{S}, \alpha} \left((\text{Tr}_{E_n} \circ \mathcal{U}_n)(\rho'_n) \left\| (\text{Tr}_{E_n} \circ \mathcal{V}_n)(\sigma'_n) \right. \right) \geq D_{\text{S}, \alpha} \left(\text{Tr}_{A_n} \rho'_n \left\| \text{Tr}_{A_n} \sigma'_n \right. \right) + D_{\text{S}, \alpha}^{\text{inf}, \infty}(\text{Tr}_{E_n} \circ \mathcal{U}_n \| \text{Tr}_{E_n} \circ \mathcal{V}_n). \quad (65)$$

Now note that $\text{Tr}_{A_n} \rho'_n = \rho_{n-1}$ where $\rho_{n-1} = \text{Tr}_{R_{n-1} E_{n-1}} \prod_{i=1}^{n-1} \mathcal{U}_i \circ \mathcal{P}^i$ and similarly for σ . As a result, applying the chain rule $n-1$ times, we get

$$D_{\text{S}, \alpha} \left(\rho_n \left\| \sigma_n \right. \right) \geq \sum_{i=1}^n D_{\text{S}, \alpha}^{\text{inf}, \infty}(\text{Tr}_{E_i} \circ \mathcal{U}_i \| \text{Tr}_{E_i} \circ \mathcal{V}_i) \quad (66)$$

$$= \sum_{i=1}^n D_{\text{M}, \alpha}^{\text{inf}, \infty}(\text{Tr}_{E_i} \circ \mathcal{U}_i \| \text{Tr}_{E_i} \circ \mathcal{V}_i) \quad (67)$$

$$\geq \sum_{i=1}^n \frac{1}{m} D_{\text{M}, \alpha}^{\text{inf}}((\text{Tr}_{E_i} \circ \mathcal{U}_i)^{\otimes m} \| (\text{Tr}_{E_i} \circ \mathcal{V}_i)^{\otimes m}) \quad (68)$$

where $m \geq 2$ and we used [41, Lemma 28] for the equality and the superadditivity of $D_{\text{M}, \alpha}$ in [41, Lemma 21] for the last inequality. Note that both of these results were applied for the family of states $\mathcal{A}_m'' = (\text{Tr}_{E_i} \circ \mathcal{U}_i)^{\otimes m}(\mathcal{D})$ and $\mathcal{B}_m'' = (\text{Tr}_{E_i} \circ \mathcal{V}_i)^{\otimes m}(\mathcal{D})$ which satisfies [41, Assumption 24], as shown in the proof of the adversarial quantum Stein's lemma.

Observe that assumption (62) implies assumption (*) in [41, Lemma 30] as $\log \text{Tr}(\mathcal{V}_i^{\otimes m}(\sigma)) \leq m \log \lambda_{\max}(\mathcal{V}_i^\dagger(I)) \leq \frac{C}{4} m$. As a result, [41, Lemma 30] gives

$$\begin{aligned} & \frac{1}{m} D_{\text{M}, \alpha}^{\text{inf}}((\text{Tr}_{E_i} \circ \mathcal{U}_i)^{\otimes m} \| (\text{Tr}_{E_i} \circ \mathcal{V}_i)^{\otimes m}) \\ & \geq D^{\text{inf}, \infty}(\text{Tr}_{E_i} \circ \mathcal{U}_i \| \text{Tr}_{E_i} \circ \mathcal{V}_i) - (1 - \alpha)(2 + C)^2 m - \frac{2(d^2 + d) \log(m + d)}{m}, \end{aligned} \quad (69)$$

for $1 - \frac{1}{(2+C)m} < \alpha < 1$. Let us now choose $1 - \alpha = \frac{8d^2 \log m}{(2+C)^2 m^2}$ and assume that $m \geq \max \left(d, \left(\frac{16d^2}{2+C} \right)^2 \right)$ so that the condition $\alpha \geq 1 - \frac{1}{(2+C)m}$ is satisfied and $\log(m + d) \leq 2 \log m$.

Putting everything together, we get

$$D_{H,\varepsilon} \left(\rho_n \left\| \sigma_n \right. \right) \geq \sum_{i=1}^n D^{\text{inf},\infty}(\text{Tr}_{E_i} \circ \mathcal{U}_i \parallel \text{Tr}_{E_i} \circ \mathcal{V}_i) - n \frac{16d^2 \log m}{m} - \frac{(2+C)^2 m^2}{8d^2 \log m} \log \frac{1}{\varepsilon}. \quad (70)$$

We now choose $m = \left(\frac{64d^4 n}{(2+C)^2 \log \frac{1}{\varepsilon}} \right)^{1/3}$. With this choice

$$D_{H,\varepsilon} \left(\rho_n \left\| \sigma_n \right. \right) \geq \sum_{i=1}^n D^{\text{inf},\infty}(\text{Tr}_{E_i} \circ \mathcal{U}_i \parallel \text{Tr}_{E_i} \circ \mathcal{V}_i) - C' n^{2/3} \log n \log^{1/3} \frac{1}{\varepsilon}, \quad (71)$$

for a constant C' that only depends on C and d . \square




FIG. 6. Illustration of the setting of the entropy accumulation.

As a corollary, we get a weak form of the entropy accumulation statement for the smoothed max-entropy obtained in [51]. We recall that the smoothed max-entropy is defined as

$$H_{\max}^{\varepsilon}(B|C)_{\rho} = \log \inf_{\substack{\tilde{\rho}_{BC} \in \mathcal{H}_{+}(BC) \\ \text{Tr}(\tilde{\rho}) \leq 1 \\ P(\rho, \tilde{\rho}) \leq \varepsilon}} \sup_{\sigma_C \in \mathcal{D}(C)} \left\| \tilde{\rho}_{BC}^{\frac{1}{2}} I_B \otimes \sigma_C^{\frac{1}{2}} \right\|_1^2, \quad (72)$$

where $P(\rho, \sigma) = \sqrt{1 - F(\rho, \sigma)^2}$ is the purified distance with $F(\rho, \sigma) := \|\sqrt{\rho}\sqrt{\sigma}\|_1 + \sqrt{(1 - \text{Tr} \rho)(1 - \text{Tr} \sigma)}$.

Corollary 8 (*H_{max}-entropy accumulation.*) *Let $\mathcal{N}'_i \in \text{CPTP}(Y_{i-1}:S_i C_i Y_i)$ be quantum channels and $\rho_{Y_0} \in \mathcal{D}(Y_0)$ be a quantum state. Define the state $\rho_{S_1 \dots S_n C_1 \dots C_n Y_n} = \prod_{i=1}^n \mathcal{N}'_i(\rho_{Y_0})$. We have for $\varepsilon \in [0, 1/2]$,*

$$H_{\max}^{\varepsilon}(S_1 \dots S_n | C_1 \dots C_n)_{\rho} \leq \sum_{i=1}^n \sup_{\omega \in \mathcal{D}(Y_{i-1})} H(S_i | C_i)_{\mathcal{N}'_i(\omega)} + K n^{2/3} \log n \log^{1/3} \frac{1}{\varepsilon}, \quad (73)$$

where K is a constant only depending on $\max_i \dim S_i C_i$.

Proof. We apply the relative entropy accumulation theorem as stated in the main text with the following replacements: letting C'_i be isomorphic to C_i , we set $A_{i+1} \leftarrow Y_i C'_{i+1} \dots C'_n$ for $i = 0$ to $n - 1$, $A_{n+1} \leftarrow Y_n$, $B_i \leftarrow S_i C'_i$ for $i = 1$ to n , $\mathcal{N}'_i \leftarrow \mathcal{N}'_i \otimes \text{Tr}_{C'_i} \otimes \text{id}_{C'_{i+1} \dots C'_n}$ (here, id refers to the identity map) and finally \mathcal{M}_i is defined as $\mathcal{M}_i(X_{Y_{i-1} C'_i \dots C'_n}) = I_{S_i} \otimes X_{C_i C'_{i+1} \dots C'_n} \otimes |0\rangle\langle 0|_{Y_i}$, where $|0\rangle\langle 0|_{Y_i}$ is an arbitrary fixed state in $\mathcal{D}(Y_i)$. An illustrative figure is given in Figure 6.

Theorem 7 gives for the left hand side of (63) and any $\sigma \in \mathcal{D}(C'_1 \dots C'_n)$:

$$\begin{aligned} D_{\text{H},\varepsilon} & \left(\text{Tr}_{A_{n+1}} \circ \prod_{i=1}^n \mathcal{N}_i(\rho_{Y_0} \otimes \sigma_{C'_1 \dots C'_n}) \left\| \text{Tr}_{A_{n+1}} \circ \prod_{i=1}^n \mathcal{M}_i(|0\rangle\langle 0|_{Y_0} \otimes \sigma_{C'_1 \dots C'_n}) \right. \right) \\ & = D_{\text{H},\varepsilon}(\rho_{S_1 \dots S_n C_1 \dots C_n} \| I_{S_1 \dots S_n} \otimes \sigma_{C_1 \dots C_n}). \end{aligned} \quad (74)$$

As the right hand side of (63) does not depend on σ , we can take an infimum over σ and use Proposition 10 to get the following

$$\begin{aligned} \inf_{\sigma} D_{\text{H},\varepsilon} & \left(\text{Tr}_{A_{n+1}} \circ \prod_{i=1}^n \mathcal{N}_i(\rho_{Y_0} \otimes \sigma_{C'_1 \dots C'_n}) \left\| \text{Tr}_{A_{n+1}} \circ \prod_{i=1}^n \mathcal{M}_i(|0\rangle\langle 0|_{Y_0} \otimes \sigma_{C'_1 \dots C'_n}) \right. \right) \\ & \leq -H_{\text{max}}^{\sqrt{2\varepsilon}}(S_1 \dots S_n | C_1 \dots C_n)_{\rho}. \end{aligned} \quad (75)$$

On the right hand side of (63), we have terms of the form $D^{\text{inf},\infty}(\text{Tr}_{A_{i+1}} \circ \mathcal{N}'_i \| \text{Tr}_{A_{i+1}} \circ \mathcal{M}_i)$. Note that for $\omega \in \mathcal{D}(Y_{i-1} C'_i \dots C'_n)$, we have $(\text{Tr}_{A_{i+1}} \circ \mathcal{N}'_i)(\omega) = \text{Tr}_{Y_i} \mathcal{N}'_i(\omega_{Y_{i-1}})$ and $(\text{Tr}_{A_{i+1}} \circ \mathcal{M}_i)(\omega) = I_{S_i} \otimes \omega_{C_i}$. As a result, $D^{\text{inf}}(\text{Tr}_{A_{i+1}} \circ \mathcal{N}'_i \| \text{Tr}_{A_{i+1}} \circ \mathcal{M}_i) = \inf_{\omega \in \mathcal{D}(Y_{i-1})} -H(S_i | C_i)_{\mathcal{N}'_i(\omega)}$, where we used the fact that $-H(B|C)_{\rho} = \inf_{\sigma \in \mathcal{D}(C)} D(\rho_{BC} \| I_B \otimes \sigma_C)$. We now need to evaluate the regularization:

$$\begin{aligned} D^{\text{inf},\infty}(\text{Tr}_{A_{i+1}} \circ \mathcal{N}'_i \| \text{Tr}_{A_{i+1}} \circ \mathcal{M}_i) & \\ & = \inf_{m \geq 1} \frac{1}{m} \inf_{\substack{\omega \in \mathcal{D}((Y_{i-1})^{\otimes m}) \\ \sigma \in \mathcal{D}((C_i \dots C_n)^{\otimes m})}} D((\text{Tr}_{Y_i} \circ \mathcal{N}'_i)^{\otimes m}(\omega) \| I_{S_{i,1} \dots S_{i,m}} \otimes \sigma_{C_{i,1} \dots C_{i,m}}) \end{aligned} \quad (76)$$

$$= - \sup_{m \geq 1} \sup_{\omega} \frac{1}{m} H(S_{i,1} \dots S_{i,m} | C_{i,1} \dots C_{i,m})_{\mathcal{N}'_i^{\otimes m}(\omega)} \quad (77)$$

$$= - \sup_{m \geq 1} \sup_{\omega} \frac{1}{m} \sum_{j=1}^m H(S_{i,j} | C_{i,1} \dots C_{i,m} S_{i,1} \dots S_{i,j-1})_{\mathcal{N}'_i^{\otimes m}(\omega)} \quad (78)$$

$$\geq - \sup_{m \geq 1} \sup_{\omega} \frac{1}{m} \sum_{j=1}^m H(S_{i,j} | C_{i,j})_{\mathcal{N}'_i(\omega_{Y_{i,j}})}, \quad (79)$$

where we used the chain rule and then strong subadditivity of the von Neumann entropy. Now note that each term $H(S_{i,j} | C_{i,j})_{\mathcal{N}'_i(\omega_{Y_{i,j}})} \leq \sup_{\omega} H(S_i | C_i)_{\mathcal{N}'_i(\omega)}$. As a result, $D^{\text{inf},\infty}(\text{Tr}_{A_{i+1}} \circ \mathcal{N}'_i \| \text{Tr}_{A_{i+1}} \circ \mathcal{M}_i) \geq - \sup_{\omega \in \mathcal{D}(Y_{i-1})} H(S_i | C_i)_{\mathcal{N}'_i(\omega)}$ which shows that in this case, the regularization is not needed. In order to apply Theorem 7, we need to check condition (62). First, we have for any $\omega \in \mathcal{D}(Y_i)$, $\text{Tr}(\mathcal{M}_i(\omega)) = \dim S_i$. In addition, for $\alpha \in [1/2, 1]$, we have

$$\begin{aligned} D_{\text{P},\alpha}^{\text{inf}}((\text{Tr}_{A_{i+1}} \circ \mathcal{N}'_i)^{\otimes m} \| (\text{Tr}_{A_{i+1}} \circ \mathcal{M}_i)^{\otimes m}) & \\ & = \inf_{\substack{\omega \in \mathcal{D}((Y_{i-1})^{\otimes m}) \\ \sigma \in \mathcal{D}((C_i \dots C_n)^{\otimes m})}} D_{\text{P},\alpha}((\text{Tr}_{Y_i} \circ \mathcal{N}'_i)^{\otimes m}(\omega_{Y_{i-1,1} \dots Y_{i-1,m}}) \| I_{S_{i,1} \dots S_{i,m}} \otimes \sigma_{C_{i,1} \dots C_{i,m}}) \end{aligned} \quad (80)$$

$$= - \sup_{\omega} H_{\text{P},\alpha}^{\uparrow}(S_{i,1} \dots S_{i,m} | C_{i,1} \dots C_{i,m})_{\mathcal{N}'_i^{\otimes m}(\omega)}, \quad (81)$$

using the notation $H_{\text{P},\alpha}^{\uparrow}(D|E)_{\rho} = - \inf_{\sigma \in \mathcal{D}(E)} D_{\text{P},\alpha}(\rho_{DE} \| I_D \otimes \sigma_E)$ from [59]. It is shown in [59, Lemma 1], that an explicit choice of σ achieves this infimum namely $\sigma_E^{(\alpha)} = \frac{(\text{Tr}_D \rho_{DE}^{\alpha})^{1/\alpha}}{\text{Tr}(\text{Tr}_D \rho_{DE}^{\alpha})^{1/\alpha}}$. But in Lemma 9, we showed that for this choice $D_{\text{P},3/2}(\rho_{DE} \| I_D \otimes \sigma_E^{(\alpha)}) \leq 4 \log \dim D$. Applying this to the state $\rho = (\text{Tr}_{A_{i+1}} \circ \mathcal{N}'_i)^{\otimes m}(\omega)$, an optimal choice for σ in (80) is given by $\sigma_m^{(\alpha)} = \frac{(\text{Tr}_{B_i^{\otimes m}} \rho^{\alpha})^{1/\alpha}}{\text{Tr}(\text{Tr}_{B_i^{\otimes m}} \rho^{\alpha})^{1/\alpha}}$. We get that for any $\omega \in \mathcal{D}(R_{i-1}^{\otimes m})$,

$$D_{\text{P},3/2}((\text{Tr}_{A_{i+1}} \circ \mathcal{N}'_i)^{\otimes m}(\omega) \| I_{S_i^{\otimes m}} \otimes \sigma_m^{(\alpha)}) \leq 4m \log \dim S_i. \quad (82)$$

This means that choosing $C = 16 \max_i \log \dim S_i$ satisfies condition (62). \square

Note that the second order term we achieve with our proof technique is worse than the one achieved in [50] and [51]. In addition, the statement of [51] is stronger in that it also includes conditioning on the system Y_i provided a non-signalling assumption is satisfied. Nevertheless, we believe that our new proof technique, which is more naturally adapted to find upper bounds on the max-entropy H_{\max} (as opposed to the techniques of [51] which naturally apply to the min-entropy H_{\min}) could lead to insights and improvements for the applications of entropy accumulation. However, this is outside the scope of this paper and we leave it for future work.

USEFUL PROPERTIES

Lemma 9 *Let $\alpha \in [1/2, 1]$, $\rho_{AB} \in \mathcal{D}(AB)$ and $\sigma_B^{(\alpha)} = \frac{(\text{Tr}_A \rho_{AB}^\alpha)^{\frac{1}{\alpha}}}{Z}$, where $Z = \text{Tr} \left((\text{Tr}_A \rho_{AB}^\alpha)^{\frac{1}{\alpha}} \right)$. Let $d_A = \dim A$. Then it holds that*

$$D_{\mathbb{P}, \frac{3}{2}}(\rho_{AB} \| I_A \otimes \sigma_B^{(\alpha)}) \leq 4 \log d_A. \quad (83)$$

Note that the choice of the parameter $\frac{3}{2}$ is not arbitrary and this lemma does not hold for higher values. See the discussion in the proof of [60, Corollary III.5].

Proof. We have

$$\text{Tr} \left(\rho_{AB}^{3/2} (I_A \otimes \sigma_B^{(\alpha)})^{-1/2} \right) = Z^{1/2} \text{Tr} \left(\rho_{AB}^{3/2} I_A \otimes (\text{Tr}_A \rho_{AB}^\alpha)^{-\frac{1}{2\alpha}} \right). \quad (84)$$

We start by showing that $Z \leq d_A$. In fact, we use the operator Jensen inequality for the operator concave function $x \mapsto x^\alpha$ as follows:

$$\text{Tr}_A(\rho_{AB}^\alpha) = d_A \sum_a \frac{\langle a | \rho_{AB}^\alpha | a \rangle}{\sqrt{d_A}} \frac{|a\rangle}{\sqrt{d_A}} \quad (85)$$

$$\leq d_A \left(\sum_a \frac{\langle a | \rho_{AB} | a \rangle}{\sqrt{d_A}} \frac{|a\rangle}{\sqrt{d_A}} \right)^\alpha \quad (86)$$

$$= d_A^{1-\alpha} \rho_B^\alpha. \quad (87)$$

As a result, we get

$$Z \leq \text{Tr}((d_A^{1-\alpha} \rho_B^\alpha)^{\frac{1}{\alpha}}) = d_A^{\frac{1-\alpha}{\alpha}} \leq d_A. \quad (88)$$

For the inequality, we used the fact that the function $x \mapsto x^{1/\alpha}$ is monotone and continuous and thus $X \mapsto \text{Tr}(X^{\frac{1}{\alpha}})$ is monotone (see e.g., [61, Section 2.2]).

Now let us consider

$$\text{Tr} \left(\rho_{AB}^{3/2} I_A \otimes (\text{Tr}_A \rho_{AB}^\alpha)^{\frac{1}{\alpha}} \right) \leq \text{Tr} \left(\rho_{AB} I_A \otimes (\text{Tr}_A \rho_{AB})^{-\frac{1}{2\alpha}} \right) \quad (89)$$

$$\leq \text{Tr}(\rho_{AB} \rho_B^{-1}) \quad (90)$$

$$\leq d_A, \quad (91)$$

where we used for the first inequality the fact that $\rho_{AB}^{3/2} \leq \rho_{AB}$, $\rho_{AB}^\alpha \geq \rho_{AB}$ and the operator anti-monotonicity of the function $x \mapsto x^{-\frac{1}{2\alpha}}$, and for the second inequality the fact that $\rho_B^{-\frac{1}{2\alpha}} \geq \rho_B$. As a result,

$$D_{\mathbb{P}, \frac{3}{2}}(\rho_{AB} \| I_A \otimes \sigma_B^{(\alpha)}) \leq 2 \log(d_A^{1/2} d_A) \leq 4 \log d_A. \quad (92)$$

\square

Proposition 10 *For $0 \leq \varepsilon < \frac{1}{2}$, we have*

$$H_{\max}^{\sqrt{2\varepsilon}}(B|C)_\rho \leq - \inf_{\sigma_C \in \mathcal{D}(C)} D_{\text{H}, \varepsilon}(\rho_{BC} \| I_B \otimes \sigma_C). \quad (93)$$

Proof. Using [41, Lemma 31] with $\mathcal{A} = \{\rho_{BC}\}$ and $\mathcal{B} = \{I_B \otimes \sigma_C : \sigma_C \in \mathcal{D}(C)\}$, we can write

$$- \inf_{\sigma_C \in \mathcal{D}(C)} D_{\text{H},\varepsilon}(\rho_{BC} \| I_B \otimes \sigma_C) = \log \inf_{0 \leq M \leq I} \left\{ \sup_{\sigma_C \in \mathcal{D}(C)} \{\text{Tr}(M_{BC} I_B \otimes \sigma_C) : \text{Tr}(M_{BC} \rho_{BC}) \geq 1 - \varepsilon\} \right\}. \quad (94)$$

Let M_{BC} be such that $\text{Tr}(M_{BC} \rho_{BC}) \geq 1 - \varepsilon$, define $\tilde{\rho}_{BC} = \sqrt{M_{BC}} \rho_{BC} \sqrt{M_{BC}}$. Then, by the gentle measurement lemma, we have $P(\rho, \tilde{\rho}) \leq \sqrt{2\varepsilon}$ (see e.g., [62, Lemma A.3]). Then, using Lemma 11, we get

$$\sup_{\sigma_C \in \mathcal{D}(C)} \left\| \left(\sqrt{M_{BC}} \rho_{BC} \sqrt{M_{BC}} \right)^{\frac{1}{2}} I_B \otimes \sigma_C^{\frac{1}{2}} \right\|_1^2 \leq \sup_{\sigma_C} \text{Tr}(M_{BC} I_B \otimes \sigma_C) \quad (95)$$

As such,

$$- \inf_{\sigma_C \in \mathcal{D}(C)} D_{\text{H},\varepsilon}(\rho_{BC} \| I_B \otimes \sigma_C) \geq \log \inf_{0 \leq M \leq I} \sup_{\sigma_C} \left\{ \left\| \left(\sqrt{M_{BC}} \rho_{BC} \sqrt{M_{BC}} \right)^{\frac{1}{2}} I_B \otimes \sigma_C^{\frac{1}{2}} \right\|_1^2 : \text{Tr}(M_{BC} \rho_{BC}) \geq 1 - \varepsilon \right\} \quad (96)$$

$$\geq \log \inf_{\substack{\tilde{\rho}_{BC} \in \mathcal{H}_+(BC) \\ \text{Tr}(\tilde{\rho}) \leq 1 \\ P(\rho, \tilde{\rho}) \leq \sqrt{2\varepsilon}}} \sup_{\sigma_C \in \mathcal{D}(C)} \left\| \tilde{\rho}_{BC}^{\frac{1}{2}} I_B \otimes \sigma_C^{\frac{1}{2}} \right\|_1^2 \quad (97)$$

$$= H_{\max}^{\sqrt{2\varepsilon}}(B|C)_\rho. \quad (98)$$

□

Lemma 11 *Let $\rho \in \mathcal{D}(A)$ and $\sigma, M \in \mathcal{H}_+(A)$. Then it holds that*

$$\left\| \left(\sqrt{M} \rho \sqrt{M} \right)^{\frac{1}{2}} \sigma^{\frac{1}{2}} \right\|_1^2 \leq \text{Tr}(M \sigma). \quad (99)$$

Proof. This fact is used in [62, Proposition 4.2]. It uses the semidefinite program for the fidelity $\|\sqrt{\omega} \sqrt{\theta}\|_1^2 = \min\{\text{Tr}(Z\theta) : \omega_{AE} \leq Z \otimes I_E, Z \geq 0\}$ [63, Section 5], where ω_{AE} is a purification of ω . Let ρ_{AE} be a purification of ρ_A . Then we have $\sqrt{M_A} \rho_{AE} \sqrt{M_A} \leq M_A \otimes I_E$ and $\sqrt{M_A} \rho_{AE} \sqrt{M_A}$ is a purification of $\sqrt{M} \rho_A \sqrt{M}$. As a result $Z = M$ is feasible for the semidefinite program above and we get the desired result. □