

Polynomial optimization on the sphere and quantum entanglement testing

Kun Fang

Joint work with Hamza Fawzi

Presented at ICCOPT 2019, Berlin



UNIVERSITY OF
CAMBRIDGE

Talk Outline

- ◎ Polynomial Optimization and SOS Hierarchy
- ◎ An improved Convergence Rate
Main Result and Proof Strategy
- ◎ Relation to Entanglement Testing
SOS Hierarchy (polynomial) v.s. DPS Hierarchy (quantum)
- ◎ Summary and Discussions

Polynomial Optimization on the Sphere



Polynomial Optimization on the Sphere



Given a multivariate polynomial $p(x)$ with $x = (x_1, \dots, x_d)$

Computing the maximal value $p_{\max} = \max_{x \in S^{d-1}} p(x)$

Over the unit sphere $S^{d-1} = \{x \in \mathbb{R}^d : x_1^2 + \dots + x_d^2 = 1\}$

Applications:

- the largest stable/independent set of a graph
 - *Degree 3* polynomial opt. on the sphere (e.g. [Nesterov'03, De Klerk'08])
- $2 \rightarrow 4$ norm of a matrix A , $p(x) = \|Ax\|_4^4$
 - *Degree 4* polynomial opt. on the sphere (e.g. [Barak et al.'12])
- Best Separable State problem in quantum information theory
 - *Degree 4* polynomial opt. on the product of spheres (e.g. [Barak-Kothari-Steurer'17])
- ...

Polynomial Optimization on the Sphere



Given a multivariate polynomial $p(x)$ with $x = (x_1, \dots, x_d)$

Computing the maximal value $p_{\max} = \max_{x \in S^{d-1}} p(x)$

Over the unit sphere $S^{d-1} = \{x \in \mathbb{R}^d : x_1^2 + \dots + x_d^2 = 1\}$

Difficulty:

- Degree = 2, efficiently solved as an eigenvalue problem;
- Degree > 2, **NP-hard** in general!


Solution:

- Sum-of-square (SOS) hierarchy [Parrilo'00; Lasserre'01]
where each level is efficiently computable by semidefinite program

Sum-of-Square (SOS) Hierarchy

$$\ell\text{-SOS} = \left\{ \underset{\substack{\uparrow \\ \text{polynomial}}}{p(x)} = \sum_i \underset{\substack{\uparrow \\ \text{polynomial}}}{q_i(x)}^2 \text{ on } S^{d-1} \mid \text{s.t. } \deg(q_i) \leq \ell \right\}$$


$$\text{NN} = \{p(x) : p(x) \geq 0, \forall x \in S^{d-1}\}$$



Relation with polynomial optimization:

$$\begin{aligned} p_{\max} &= \max_{x \in S^{d-1}} p(x) = \min\{\gamma \in \mathbb{R} : \boxed{\gamma - p \in \text{NN}} \text{ on } S^{d-1}\} \\ &\leq \min\{\gamma \in \mathbb{R} : \boxed{\gamma - p \in \ell\text{-SOS}} \text{ on } S^{d-1}\} \quad \text{restriction} \\ &= p_\ell \quad [\text{SDP of size } d^{O(\ell)}] \end{aligned}$$

Approaching $p_{\max}$ from above:



Main Result: improved Convergence Rate



Q: How fast does p_l converge to $p_{\max}$?

A: [Reznick'95; Doherty-Wehner'12], convergence rate at least $O(d/\ell)$

Q: Can we further sharpen the convergence rate?

(see recent works by de Klerk & Laurent 1811.05439 & 1904.08828)

A: *Positive answer in this work*, convergence rate at least $O((d/\ell)^2)$

Quadratic improvement

Main Result (technical statement)

Suppose $p(x_1, \dots, x_d)$ is a homo. poly. of degree $2n$ in d variables with $n \leq d$,

$$1 \leq \frac{p_l - p_{\min}}{p_{\max} - p_{\min}} \leq 1 + \left(C_n \cdot \frac{d}{\ell} \right)^2 \quad \text{for all } \ell \geq C_n d$$

reference point
 $\downarrow$
 $p_{\min} = \min_{x \in S^{d-1}} p(x)$

$\downarrow$
A constant depends only on n .

A Stronger Result [take home message]

Matrix-valued polynomials:

$$x = (x_1, \dots, x_d)$$

$F(x) \in \mathbf{S}^k[x]$: k by k matrix with polynomial entries, symmetric for any x .

$F(x) \geq 0$: positive semidefinite matrix for any x .

$$F(x) \in \ell\text{-SOS} \quad \text{if} \quad F(x) = \sum_j U_j(x) U_j(x)^\top, \quad \deg(U_j) \leq \ell$$

- Remark:**
- These definitions reduce to (scalar-valued) polynomial if $k = 1$;
 - But the results cannot be trivially extended. (e.g. Nonnegative quadratic polynomial is necessarily a SOS. But not true for matrix-valued case.)

For any homo. matrix-valued poly. $F(x) \in \mathbf{S}^k[x]$ of degree $2n$ in d variables with $n \leq d$ and $0 \leq F(x) \leq I$ for all $x \in S^{d-1}$,

$$F + C'_n \left(\frac{d}{\ell} \right)^2 \cdot I \in \ell\text{-SOS on } S^{d-1} \quad \text{for all } \ell \geq C_n d$$

A small perturbation δ

Identity

A remarkable fact: the result is totally independent on the size of the matrix $F(x)$.

A Stronger Result

Main Result (technical statement)

Suppose $p(x_1, \dots, x_d)$ is a homo. poly. of degree $2n$ in d variables with $n \leq d$,

$$1 \leq \frac{p_\ell - p_{\min}}{p_{\max} - p_{\min}} \leq 1 + \left(C_n \cdot \frac{d}{\ell} \right)^2 \quad \text{for all } \ell \geq C_n d$$



$$F = \frac{p_{\max} - p}{p_{\max} - p_{\min}}$$

For any homo. matrix-valued poly. $F(x) \in \mathbf{S}^k[x]$ of degree $2n$ in d variables with $n \leq d$ and $0 \leq F(x) \leq I$ for all $x \in S^{d-1}$,

$$F + C'_n \left(\frac{d}{\ell} \right)^2 \cdot I \in \ell\text{-SOS on } S^{d-1} \quad \text{for all } \ell \geq C_n d$$

A small perturbation δ

Identity

A remarkable fact: the result is totally independent on the size of the matrix $F(x)$.

Convergence Rate

Proof Outline

Proof Outline


Goal: Given poly. $0 \leq F \leq 1$. Find $\delta > 0$ such that $\tilde{F} = F + \delta$ is ℓ -SOS

How to construct SOS:

For any polynomial: $q(t) : [-1, 1] \rightarrow \mathbb{R}$ of $\deg(q) = \ell$, consider $K(x, y) = q(\langle x, y \rangle)^2$

$$\begin{aligned} \underline{(Kh)(x)} &= \int_{y \in S^{d-1}} \underline{K(x, y)} \underline{h(y)} d\sigma(y) & \forall x \in S^{d-1} \\ &= \underbrace{\int_{y \in S^{d-1}}}_{\text{sum}} \underbrace{q(\langle x, y \rangle)^2}_{\text{Poly}^2} \underbrace{h(y) d\sigma(y)}_{\text{nonnegative weight}} & \forall x \in S^{d-1} \end{aligned}$$

Key observation: $h(y) \geq 0 \implies Kh \in \ell\text{-SOS}$ [Reznick'95; Doherty-Wehner'12 ;Parrilo'13]



$$\tilde{F} = F + \delta \text{ is } \ell\text{-SOS} \quad \leftarrow \quad \|K^{-1}\tilde{F} - \tilde{F}\|_{\infty} \leq \delta$$

Estimate $\|K^{-1}\tilde{F} - \tilde{F}\|_{\infty} \leq \delta$ Rotation invariant kernel $K(x, y) = q(\langle x, y \rangle)^2$

Some well-studied results

Harmonic Decomposition $F = F_0 + F_2 + \cdots F_{2n}$ $\tilde{F} = (F_0 + \delta) + F_2 + \cdots F_{2n}$

Kernel Decomposition $\phi = q^2 = \lambda_0 C_0 + \lambda_1 C_1 + \cdots \lambda_{2\ell} C_{2\ell}$ [C_k Gegenbauer]

Funk-Hecke formula $K\tilde{F} = \lambda_0(F_0 + \delta) + \lambda_2 F_2 + \cdots \lambda_{2n} F_{2n}$

$$K^{-1}\tilde{F} = \lambda_0^{-1}(F_0 + \delta) + \lambda_2^{-1} F_2 + \cdots \lambda_{2n}^{-1} F_{2n}$$

$$\|K^{-1}(\tilde{F}) - \tilde{F}\|_{\infty} = \left\| \sum_{k=1}^n \left(\frac{1}{\lambda_{2k}} - 1 \right) F_{2k} \right\|_{\infty} \leq \sum_{k=1}^n \underbrace{\left| \frac{1}{\lambda_{2k}} - 1 \right|}_{\text{red}} \|F_{2k}\|_{\infty}$$

○ Since $0 \leq F \leq 1$, its harmonic components won't be too large $\|F_{2k}\|_{\infty} \leq B_{2n} \|F\|_{\infty}$

○ Estimate $\sum_{k=1}^n \left| \frac{1}{\lambda_{2k}} - 1 \right|$  $\sum_{k=1}^n \left| \frac{1}{\lambda_{2k}} - 1 \right| \leq 2 \underbrace{\sum_{k=1}^n (1 - \lambda_{2k})}_{\text{red box}}$
 $\ell \geq 2nd$

$\tilde{F} = F + \delta$ is ℓ -SOS

$$\|K^{-1}\tilde{F} - \tilde{F}\|_{\infty} \leq \delta$$

$$\sum_{k=1}^n (1 - \lambda_{2k}) \leq \delta$$

Estimate $\sum_{k=1}^n (1 - \lambda_{2k}) \leq \delta$

$$\lambda_i = \frac{\omega_{d-1}}{\omega_d} \int_{-1}^1 \phi(t) \frac{C_i(t)}{C_i(1)} (1 - t^2)^{\frac{d-3}{2}} dt \quad \phi = q^2 = \lambda_0 C_0 + \lambda_1 C_1 + \cdots \lambda_{2\ell} C_{2\ell}$$

- If we choose polynomial $q(t) \propto t^{\ell}$, each coefficient can be computed explicitly. Observe that λ_i scales as $O(d/\ell)$. Recover results by [Reznick'95; Doherty-Wehner'12].
- **To obtain a better result, we do not choose specific $q(t)$ at this moment.**

$$\phi(t) = [q(t)]^2 = \left[\sum_{i=0}^{\ell} e_i \frac{C_i(t)}{\sqrt{C_i(1)}} \right]^2 \quad e = [e_0 \ e_1 \ \cdots \ e_{\ell}]^{\top}$$

$$\lambda_{2k} = e^{\top} \mathcal{T}[C_{2k}/C_{2k}(1)] e \quad \mathcal{T}[g]_{i,j} = \frac{\omega_{d-1}}{\omega_d} \int_{-1}^1 \frac{C_i(t)}{\sqrt{C_i(1)}} \frac{C_j(t)}{\sqrt{C_j(1)}} g(t) (1 - t^2)^{\frac{d-3}{2}} dt$$

Generalized Toeplitz matrix

$$\sum_{k=1}^n (1 - \lambda_{2k}) = n (1 - e^{\top} \mathcal{T}[h] e)$$




$$1 - \lambda_{\max}(\mathcal{T}[h]) \leq \delta$$

$$h = \frac{1}{n} \sum_{k=1}^n \frac{C_{2k}}{C_{2k}(1)}$$

$$\boxed{\tilde{F} = F + \delta \text{ is } \ell\text{-SOS}} \leftarrow \boxed{\|K^{-1}\tilde{F} - \tilde{F}\|_{\infty} \leq \delta} \leftarrow \boxed{\sum_{k=1}^n (1 - \lambda_{2k}) \leq \delta} \leftarrow \boxed{1 - \lambda_{\max}(\mathcal{T}[h]) \leq \delta}$$

Estimate $1 - \lambda_{\max}(\mathcal{T}[h]) \leq \delta$ $h = \frac{1}{n} \sum_{k=1}^n \frac{C_{2k}}{C_{2k}(1)}$

- For linear polynomial f , we have a good understanding of the eigenvalues of $\mathcal{T}[f]$.
- h is non-linear -> Consider the tangent line at $t = 1$, $\bar{h}(t) = h'(1)(t - 1) + h(1)$



This completes the proof.

Remark: The estimation of $\lambda_{\max}(\mathcal{T}[h])$ is tight.

$d = 4, n = 2$

Some Remarks

Recall the result

For any homo. matrix-valued poly. $F(x) \in \mathbf{S}^k[x]$ of degree $2n$ in d variables with $n \leq d$ and $0 \leq F(x) \leq I$ for all $x \in S^{d-1}$

$$F + C'_n \left(\frac{d}{\ell} \right)^2 \cdot I \in \ell\text{-SOS on } S^{d-1} \quad \text{for all } \ell \geq C_n d$$

Some Remarks:

- The proof works for *matrix-valued polynomials*.
- The proof works for polynomials on the *complex sphere*.
- We can estimate the convergence for *all values of level*, not just $\ell \geq C_n d$.

Entanglement Testing



Quantum States

Quantum state $\mathcal{S}(\mathcal{H}_A) = \left\{ \sum_i p_i x_i x_i^\dagger : p_i \geq 0, x_i \in \mathcal{H}_A \right\}$

Separable state $\mathcal{SEP}(\mathcal{H}_A \otimes \mathcal{H}_B) = \left\{ \sum_i p_i (x_i x_i^\dagger) \otimes (y_i y_i^\dagger) : p_i \geq 0, x_i \in \mathcal{H}_A, y_i \in \mathcal{H}_B \right\}$

Entangled state Any quantum state that is *not separable*

Q: Whether a given quantum state ρ_{AB} is entangled or not?

A: Doherty-Parrilo-Spedalieri (DPS) hierarchy

$$\mathcal{DPS}_\ell(\mathcal{H}_A \otimes \mathcal{H}_B) = \{ \rho_{AB} : \exists \rho_{AB_1 \dots B_\ell} \text{ s.t. (1, 2, 3) holds} \}$$

1. Reduction under partial trace:


$$\text{Tr}_{B_2 \dots B_\ell} [\rho_{AB_1 \underline{B_2 \dots B_\ell}}] = \rho_{AB}$$

2. Symmetry on B systems:

$$(I \otimes \Pi_{B_1 \dots B_\ell}) \rho_{AB_1 \dots \underline{B_\ell}} (I \otimes \Pi_{B_1 \dots B_\ell}) = \rho_{AB_1 \dots B_\ell}$$

3. Positive partial transpose (PPT):

$$(I_A \otimes \boxed{\mathsf{T}_{B_1} \otimes \dots \otimes \mathsf{T}_{B_s}} \otimes I_{B_{s+1}} \otimes I_{B_\ell})(\rho_{AB_1 \dots B_\ell}) \geq 0$$



Form a complete hierarchy [Doherty-Parrilo-Spedalieri'02&04]

Quantum States

Quantum state $\mathcal{S}(\mathcal{H}_A) = \left\{ \sum_i p_i x_i x_i^\dagger : p_i \geq 0, x_i \in \mathcal{H}_A \right\}$

Separable state $\mathcal{SEP}(\mathcal{H}_A \otimes \mathcal{H}_B) = \left\{ \sum_i p_i (x_i x_i^\dagger) \otimes (y_i y_i^\dagger) : p_i \geq 0, x_i \in \mathcal{H}_A, y_i \in \mathcal{H}_B \right\}$

Entangled state Any quantum state that is *not separable*

Q: Whether a given quantum state ρ_{AB} is entangled or not?

A: Doherty-Parrilo-Spedalieri (DPS) hierarchy

$$\mathcal{DPS}_\ell(\mathcal{H}_A \otimes \mathcal{H}_B) = \{ \rho_{AB} : \exists \rho_{AB_1 \dots B_\ell} \text{ s.t. (1, 2, 3) holds} \}$$

1. Reduction under partial trace:


$$\text{Tr}_{B_2 \dots B_\ell} [\rho_{AB_1 B_2 \dots B_\ell}] = \rho_{AB}$$

2. Symmetry on B systems:

$$(I \otimes \Pi_{B_1 \dots B_\ell}) \rho_{AB_1 \dots B_\ell} (I \otimes \Pi_{B_1 \dots B_\ell}) = \rho_{AB_1 \dots B_\ell}$$

3. Positive partial transpose (PPT):


$$(I_A \otimes \boxed{\mathsf{T}_{B_1} \otimes \dots \otimes \mathsf{T}_{B_s}} \otimes I_{B_{s+1}} \otimes I_{B_\ell}) (\rho_{AB_1 \dots B_\ell}) \geq 0$$



Without PPT conditions $\mathcal{EXT}_\ell(\mathcal{H}_A \otimes \mathcal{H}_B) = \{ \rho_{AB} : \exists \rho_{AB_1 \dots B_\ell} \text{ s.t. (1, 2) holds} \}$

This is a weaker hierarchy but it is still complete.

Duality Relation



For any Hermitian operator M on $\mathcal{H}_A \otimes \mathcal{H}_B$, define its associated Hermitian polynomial

$$p_M(x, \bar{x}, y, \bar{y}) = (x \otimes y)^\dagger M (x \otimes y) = \sum_{i,j,k,l} M_{ij,kl} x_i \bar{x}_k y_j \bar{y}_l \quad \forall x \in \mathbb{C}^{d_A}, y \in \mathbb{C}^{d_B}$$

Duality relation

$$\mathcal{SEP}^* = \{M \in \text{Herm}(\mathcal{H}_A \otimes \mathcal{H}_B) : p_M \text{ is nonnegative}\}$$

$$\mathcal{DPS}_\ell^* = \left\{ M \in \text{Herm}(\mathcal{H}_A \otimes \mathcal{H}_B) : \|y\|^{2(\ell-1)} p_M \text{ is rSOS} \right\} \cdots \rightarrow \sum_i q_i(x, \bar{x}, y, \bar{y})^2$$

$$\mathcal{EXT}_\ell^* = \left\{ M \in \text{Herm}(\mathcal{H}_A \otimes \mathcal{H}_B) : \|y\|^{2(\ell-1)} p_M \text{ is cSOS} \right\} \cdots \rightarrow \sum_i |g_i(x, y)|^2$$

PPT conditions determine the choice of monomials in the SOS decomposition.

Relation with [NOP'09]

[Navascues-Owari-Plenio'09]

For any quantum state $\rho_{AB} \in \text{DPS}_\ell$ with reduced state $\rho_A = \text{Tr}_B[\rho_{AB}]$

$$(1-t)\rho_{AB} + t\rho_A \otimes \frac{I_B}{d_B} \text{ is separable with } t = O\left(\frac{d_B^2}{\ell^2}\right)$$



A small perturbation

Utilizing the duality between DPS and SOS, this is equivalent to our result of matrix-valued polynomial with **degree 2**.

Recall our result in this work

For any homo. matrix-valued poly. $F(x) \in \mathbf{S}^k[x]$ of degree $2n$ in d variables with $n \leq d$ and $0 \leq F(x) \leq I$ for all $x \in S^{d-1}$

$$F + C'_n \left(\frac{d}{\ell}\right)^2 \cdot I \in \ell\text{-SOS on } S^{d-1} \quad \text{for all } \ell \geq C_n d$$




A small perturbation

Summary & Discussions

Summary

- **An quadratic improvement of convergence rate of the SOS hierarchy**
 - Works for matrix-valued polynomials
 - Works for complex variables
 - Works for all values of the level
 - **Exact Duality relation between SOS and DPS hierarchies**
 - **Connection with [Navascues-Owari-Plenio'09] from quantum community**
-

Other related works:



- Lasserre hierarchy appr. from below
[de Klerk-Laurent 1904.08828]

SOS hierarchy appr. from above
[Fang-Fawzi-This work]

Empirically much faster

- Analysis of the SOS hierarchy from the computer science community
(e.g. [Bhattiprolu et al.'17; Barak-Kothari-Steurer'17])

Question: Further sharpening the convergence rate? New techniques are required.

Thanks for your attention!

Full paper will be online soon.